

Goodness-of-Fit Tests for Random Partitions via Symmetric Polynomials

Chao Gao

CHAOGAO@GALTON.UCHICAGO.EDU

*Department of Statistics
University of Chicago
Chicago, IL 60637, USA*

Editor: Animashree Anandkumar

Abstract

We consider goodness-of-fit tests with i.i.d. samples generated from a categorical distribution (p_1, \dots, p_k) . For a given (q_1, \dots, q_k) , we test the null hypothesis whether $p_j = q_{\pi(j)}$ for some label permutation π . The uncertainty of label permutation implies that the null hypothesis is composite instead of being singular. In this paper, we construct a testing procedure using statistics that are defined as indefinite integrals of some symmetric polynomials. This method is aimed directly at the invariance of the problem, and avoids the need of matching the unknown labels. The asymptotic distribution of the testing statistic is shown to be chi-squared, and its power is proved to be nearly optimal under a local alternative hypothesis. Various degenerate structures of the null hypothesis are carefully analyzed in the paper. A two-sample version of the test is also studied.

Keywords: hypothesis testing, elementary symmetric polynomials, Lagrange interpolating polynomials, Vandermonde matrix, minimax optimality

1. Introduction

Consider a categorical distribution parameterized by (p_1, \dots, p_k) . We have i.i.d. observations X_1, \dots, X_n that follow $\mathbb{P}(X_i = j) = p_j$. A classical goodness-of-fit testing problem is to test whether or not $p_j = q_j$ for $j \in [k]$, where q_1, \dots, q_k are some given numbers. One solution is given by the famous Pearson's chi-squared test (Pearson, 1900). In this traditional formulation, it is assumed that the labels $(1, \dots, k)$ of (p_1, \dots, p_k) correspond to those of (q_1, \dots, q_k) , so that p_j can be directly compared with q_j for each $j \in [k]$. However, this assumption is not satisfied in some interesting applications. We give three examples below:

1. *Clustering models.* In a typical probabilistic setting of cluster analysis, the event $\{X_i = j\}$ means that the i th item belongs to the j th cluster, and p_j is the population frequency of the j th cluster. Here, the cluster label j does not carry any real meaning, and is present only for notational convenience. In a cluster analysis setting, the underlying object of interest is the partition of the n items instead of the cluster labels. In other words, what really matters to statisticians is the value of $\mathbb{I}\{X_i = X_{i'}\}$ (the indicator function of the event) for every pair $i \neq i'$. Therefore, a clustering model with population frequency (p_1, \dots, p_k) is equivalent to that with $(p_{\pi(1)}, \dots, p_{\pi(k)})$ with some permutation π .

2. *Word frequency analysis.* Consider two text corpora of two different languages. The word frequencies are denoted by (p_1, \dots, p_k) and (q_1, \dots, q_k) , respectively. An interesting problem in comparative linguistics is to study whether the two languages share common features by comparing (p_1, \dots, p_k) with (q_1, \dots, q_k) . For languages that are not necessarily etymologically related, the correspondence between words of the two languages are usually unclear or unknown. Therefore, a reasonable comparison of word frequencies between two languages can be conducted through comparing (p_1, \dots, p_k) with a reordered vector $(q_{\pi(1)}, \dots, q_{\pi(k)})$ for some permutation π .
3. *Simple substitution cipher.* In cryptography, a simple substitution cypher changes every character in a message to a different character systematically. Let $\{1, \dots, k\}$ be a finite alphabet of characters, and (Y_1, \dots, Y_n) denote a message to be encrypted. A simple substitution cypher is defined by a permutation σ on the alphabet $\{1, \dots, k\}$. This results in the encrypted message (X_1, \dots, X_n) with $X_i = \sigma(Y_i)$ for each $i \in [n]$. Suppose each Y_i is independently distributed by $\mathbb{P}(Y_i = j) = q_j$. Then, each X_i independently follows $\mathbb{P}(X_i = j) = p_j$, where $p_j = q_{\pi(j)}$ with $\pi = \sigma^{-1}$. If only the encrypted message is observed, inference of the probability vector (q_1, \dots, q_k) is only possible up to an unknown permutation.

Inspired by the above examples, in this paper, we consider a twist of the traditional formulation of the hypothesis testing problem. We consider the following null hypothesis:

$$H_0 : p_j = q_{\pi(j)}, \quad \text{for some } \pi \in S_k, \quad (1)$$

where (q_1, \dots, q_k) is a known vector and S_k is the set of all permutations of $[k]$. This null hypothesis implies that the labels $1, \dots, k$ do not have any meaning. For example, the vectors $(\frac{1}{2}, \frac{1}{4}, \frac{1}{4})$ and $(\frac{1}{4}, \frac{1}{4}, \frac{1}{2})$ are considered equivalent. Given i.i.d. observations X_1, \dots, X_n , one can immediately define summary statistics $n_j = \sum_{i=1}^n \mathbb{I}\{X_i = j\}$ for $j \in [k]$, which are sufficient. Since the labels of n_1, \dots, n_k are irrelevant, these sufficient statistics result in a random partition of the integer n . There are two ways to code such a random partition (Pitman, 1995): (i) by the order statistics $n_{(1)} \geq n_{(2)} \geq \dots \geq n_{(k)}$; (ii) by the numbers of terms of various sizes $m_l = \sum_{j=1}^k \mathbb{I}\{n_j = l\}$ for $l \in [n]$. It is easy to see that $\sum_{l=1}^n m_l = k$ and $\sum_{l=1}^n l m_l = n$. These two representations are equivalent because one can be derived from the other.

Inference of the probability vector (p_1, \dots, p_k) up to a label permutation using random partitions have been extensively studied in Bayesian statistics. The problem serves as a foundation for random partition models, cluster analysis and species distribution modeling. Priors that induce various exchangeable properties have been developed for the equivalent class $\{(p_{\pi(1)}, \dots, p_{\pi(k)}) : \pi \in S_k\}$. See Ewens (1972); Pitman (1995); Müller et al. (2013); Crane (2016); Gnedin and Pitman (2006); De Blasi et al. (2015); Kingman (1975); Pitman (1996) and references therein. In this paper, we take a frequentist point of view that is complementary to the Bayesian literature, and we do not treat the equivalent class $\{(p_{\pi(1)}, \dots, p_{\pi(k)}) : \pi \in S_k\}$ as random. The theory of hypothesis testing is developed within a frequentist decision-theoretic framework.

With the unknown permutation π in the null hypothesis, the classical chi-squared test by Pearson does not work anymore. Our idea of the test is based on the following class of

statistics:

$$\left\{ \sum_{j=1}^k f(n_j) : f \in \mathcal{F} \right\}, \quad (2)$$

where \mathcal{F} is the class of all measurable functions. For each $f \in \mathcal{F}$, the distribution of $\sum_{j=1}^k f(n_j)$ is identical for $p_j = q_{\pi(j)}$ with any $\pi \in S_k$. This is because (2) is a class of statistics that are invariant to the label permutation π . That is, $\sum_{j=1}^k f(n_j) = \sum_{j=1}^k f(n_{\pi(j)})$ for any $\pi \in S_k$. Moreover, it is easy to see that these statistics are all functions of the random partition because $\sum_{j=1}^k f(n_j) = \sum_{j=1}^k f(n_{(j)})$.

Choosing an appropriate class of f 's is important. We propose to use k functions f_1, \dots, f_k that satisfy the *identifiability* and the *orthogonality* conditions. The identifiability condition requires that the k equations $\sum_{j=1}^k f_l(p_j) = \sum_{j=1}^k f_l(q_j)$ for $l \in [k]$ hold if and only if $p_j = q_{\pi(j)}$ for some $\pi \in S_k$. With this condition, testing whether the null hypothesis holds is equivalent to testing whether the k equations hold. The orthogonality condition requires that the k vectors $(f_l'(q_1), \dots, f_l'(q_k))^T$ for $l \in [k]$ are orthogonal to each other. Intuitively speaking, this condition ensures that the information carried by the k statistics $\sum_{j=1}^k f_l(n_j)$ for $l \in [k]$ are mutually exclusive, which is a key ingredient that leads to optimal power under a local alternative.

In this paper, we choose f_1, \dots, f_k to be indefinite integrals of Lagrange interpolation polynomials. The choice of these polynomials satisfies the above-mentioned *identifiability* and *orthogonality* conditions. We prove that the testing statistic constructed from the k functions is asymptotically distributed by a chi-squared distribution. Moreover, we show that the power of the test is nearly optimal under a local alternative hypothesis within a decision-theoretic framework.

Our approach that uses symmetric polynomials bypasses the problem of unknown permutation π . It falls into the general umbrella of methods of moments, which are commonly used for problems that impose equivalence relations to the signals through the action of a group of transformations. For example, various method-of-moments techniques have been applied to problems including Gaussian mixture models (Hsu and Kakade, 2013), mixed membership models (Anandkumar et al., 2013), dictionary learning (Arora et al., 2014), topic models (Anandkumar et al., 2012; Arabshahi and Anandkumar, 2016) and multi-reference alignment (Perry et al., 2017). Recently, this idea was also applied to the problems of network testing by Gao and Lafferty (2017); Banerjee and Ma (2017), where the group action there is row and column permutations of the adjacency matrix of a random network.

The rest of the paper is organized as follows. In Section 2, we introduce definitions of some useful symmetric polynomials and the related Vandermonde matrix. Before getting into the testing problem for random partitions, we first solve an easier version of the problem with Gaussian observations in Section 3 and Section 4. The test using random partitions is given in Section 5. The optimality of our test is discussed in Section 6. In Section 8, we consider a two-sample version of the problem. Numerical experiments of the proposed testing procedures are given in Section 7. Finally, Section 9 is a discussion section, where we briefly analyze the property of the test on the boundary of degeneracy and discuss some open problems. The proofs of all results in the paper are given in Section 10.

We close this section by introducing the notation used in the paper. For $a, b \in \mathbb{R}$, let $a \vee b = \max(a, b)$ and $a \wedge b = \min(a, b)$. For an integer m , $[m]$ denotes the set $\{1, 2, \dots, m\}$.

Given a set S , $|S|$ denotes its cardinality, and \mathbb{I}_S is the associated indicator function. We use \mathbb{P} and \mathbb{E} to denote generic probability and expectation whose distribution is determined from the context. The noncentral chi-squared distribution with degrees of freedom k and noncentrality parameter δ^2 is denoted as χ_{k,δ^2}^2 . We will also use χ_{k,δ^2}^2 for the associated random variables.

2. Symmetric Polynomials and Vandermonde Matrix

Define a polynomial with roots $\mu_1, \dots, \mu_k \in \mathbb{R}$ by

$$f(t) = \prod_{j=1}^k (t - \mu_j).$$

It can be organized as

$$f(t) = \sum_{j=0}^k (-1)^{k-j} e_{k-j}(\mu_1, \dots, \mu_k) t^j. \quad (3)$$

The coefficient before t^j is $(-1)^{k-j} e_{k-j}(\mu_1, \dots, \mu_k)$, and $e_{k-j}(\mu_1, \dots, \mu_k)$ is called the elementary symmetric polynomial. For $l \in \{1, \dots, k\}$, the l th elementary symmetric polynomial is

$$e_l(\mu_1, \dots, \mu_k) = \sum_{1 \leq j_1 < \dots < j_l \leq k} \mu_{j_1} \cdots \mu_{j_l}.$$

When $l = 0$, we use the convention $e_0(\mu_1, \dots, \mu_k) = 1$.

The elementary symmetric polynomials can be efficiently calculated through Newton's identities. Define the l th power sum

$$p_l(\mu_1, \dots, \mu_k) = \sum_{j=1}^k \mu_j^l. \quad (4)$$

Newton's identities can be summarized through the formula

$$e_l(\mu_1, \dots, \mu_k) = \frac{1}{l} \sum_{j=1}^l (-1)^{j-1} e_{l-j}(\mu_1, \dots, \mu_k) p_j(\mu_1, \dots, \mu_k), \quad (5)$$

for $l = 1, \dots, k$.

Finally, we introduce an interesting relation between elementary symmetric polynomials and Vandermonde matrix (see Chapter 0.9.11 of Horn and Johnson (2012)). Given μ_1, \dots, μ_k that take k distinct values, define a matrix $E(\mu_1, \dots, \mu_k) \in \mathbb{R}^{k \times k}$, whose (j, l) th entry is

$$(-1)^{j-1} \frac{e_{k-j}(\mu_1, \dots, \mu_{l-1}, \mu_{l+1}, \dots, \mu_k)}{\prod_{j \in [k] \setminus \{l\}} (\mu_j - \mu_l)}. \quad (6)$$

The Vandermonde matrix $V(\mu_1, \dots, \mu_k) \in \mathbb{R}^{k \times k}$ has μ_j^{l-1} on its (j, l) th entry. Interestingly, we have

$$E(\mu_1, \dots, \mu_k) V(\mu_1, \dots, \mu_k) = V(\mu_1, \dots, \mu_k) E(\mu_1, \dots, \mu_k) = I_k. \quad (7)$$

This relation implies a formula for the determinant of $E(\mu_1, \dots, \mu_k)$:

$$\det(E(\mu_1, \dots, \mu_k)) = \frac{1}{\det(V(\mu_1, \dots, \mu_k))} = \frac{1}{\prod_{1 \leq j < l \leq k} (\mu_l - \mu_j)}. \quad (8)$$

3. The Gaussian Case

Before working with categorical distributions, we first study data generated from a Gaussian distribution. This allows us to grasp the mathematical essence of the problem without dealing with the dependence and heteroskedasticity of categorical distributions. We consider a Gaussian random vector $X \sim N(\theta, n^{-1}I_k)$. The mean vector $\theta \in \mathbb{R}^k$ consists of k numbers $\theta_1, \dots, \theta_k$. Throughout the paper, we assume $k \geq 2$ and it is a constant that does not vary with n . We would like to test whether the k numbers are identical to μ_1, \dots, μ_k after some permutation of labels. To be rigorous, introduce a distance between two vectors θ and μ ,

$$\ell(\theta, \mu) = \min_{\pi \in S_k} \sqrt{\sum_{j=1}^k (\theta_j - \mu_{\pi(j)})^2},$$

where S_k is the set of all permutations on $[k]$. Then, the hypothesis testing problem is

$$H_0 : \ell(\theta, \mu) = 0, \quad H_1 : \ell(\theta, \mu) > 0.$$

Throughout this section, we assume $\min_{j \neq l} |\mu_j - \mu_l| > 0$. The case $\min_{j \neq l} |\mu_j - \mu_l| = 0$ is degenerate and will be studied in the next section.

We use the notation μ_π to denote a k -dimensional vector whose j th entry is $\mu_{\pi(j)}$. Then, the null hypothesis can also be written as

$$\theta \in \{\mu_\pi : \pi \in S_k\}.$$

In other words, there is an equivalent class of probability distributions $\{N(\mu_\pi, n^{-1}I_k) : \pi \in S_k\}$. Thus, it is natural to consider summary statistics whose distributions are invariant under this equivalent class. This leads to the class of summary statistics

$$\left\{ \sum_{j=1}^k f(X_j) : f \in \mathcal{F} \right\},$$

where \mathcal{F} is the set of all measurable functions. For $X \sim N(\theta, n^{-1}I_k)$, it is easy to see that the distribution of $\sum_{j=1}^k f(X_j)$ only depends on the equivalent class $\{N(\theta_k, n^{-1}I_k) : \pi \in S_k\}$. This fact holds for an arbitrary $f \in \mathcal{F}$.

Since the degree of freedom of the null hypothesis is k , our strategy is to construct a testing procedure based on $\left\{ \sum_{j=1}^k f_l(X_j) : l \in [k] \right\}$. In other words, we need to choose the k functions $f_1(\cdot), \dots, f_k(\cdot)$. The following two conditions are proposed:

1. **Identifiability.** Assume $\min_{j \neq l} |\mu_j - \mu_l| > 0$. Then the equations

$$\sum_{j=1}^k f_l(\theta_j) = \sum_{j=1}^k f_l(\mu_j), \quad l = 1, \dots, k, \quad (9)$$

hold, if and only if $\ell(\theta, \mu) = 0$.

2. **Orthogonality.** Assume $\min_{j \neq l} |\mu_j - \mu_l| > 0$. Then for any $l, h \in [k]$,

$$\sum_{j=1}^k f'_l(\mu_j) f'_h(\mu_j) = \begin{cases} 1, & l = h, \\ 0, & l \neq h. \end{cases}$$

We give a few remarks regarding the two conditions. The first condition of identifiability is natural. It is required by the structure of the problem, and is necessary for the test to have power under the alternative hypothesis. The second condition implies information independence among the k summary statistics.

The k functions we propose to satisfy the two conditions are

$$f_l(t) = \frac{\int \prod_{j \in [k] \setminus \{l\}} (t - \mu_j)}{\prod_{j \in [k] \setminus \{l\}} (\mu_l - \mu_j)}, \quad l = 1, \dots, k. \quad (10)$$

The derivatives $f'_l(t) = \frac{\prod_{j \in [k] \setminus \{l\}} (t - \mu_j)}{\prod_{j \in [k] \setminus \{l\}} (\mu_l - \mu_j)}$ are called Lagrange interpolating polynomials, and it is easy to check that the second condition of orthogonality holds. Now we check the first condition of identifiability. By (3), we have

$$\prod_{j \in [k] \setminus \{l\}} (t - \mu_j) = \sum_{j=0}^{k-1} (-1)^{k-1-j} e_{k-1-j}(\mu_1, \dots, \mu_{l-1}, \mu_{l+1}, \dots, \mu_k) t^j.$$

This implies

$$f_l(t) = \sum_{j=1}^k (-1)^{k-j} \frac{e_{k-j}(\mu_1, \dots, \mu_{l-1}, \mu_{l+1}, \dots, \mu_k) t^j}{\prod_{j \in [k] \setminus \{l\}} (\mu_l - \mu_j)} \frac{1}{j}. \quad (11)$$

Therefore, the equations (9) can be written as

$$\sum_{j=1}^k (-1)^{k-j} \frac{e_{k-j}(\mu_1, \dots, \mu_{l-1}, \mu_{l+1}, \dots, \mu_k)}{\prod_{j \in [k] \setminus \{l\}} (\mu_l - \mu_j)} \Delta_j = 0, \quad l = 1, \dots, k.$$

where $\Delta_j = \frac{1}{j} \sum_{h=1}^k \theta_h^j - \frac{1}{j} \sum_{h=1}^k \mu_h^j$. In view of the definition of the matrix $E(\mu_1, \dots, \mu_k)$ in (6), we have a compact organization of the equations

$$E(\mu_1, \dots, \mu_k) \Delta = 0.$$

When the assumption $\min_{j \neq l} |\mu_j - \mu_l| > 0$ holds, the matrix $E(\mu_1, \dots, \mu_k)$ has full rank and is invertible according to (7) and (8), which immediately implies $\Delta = 0$. Equivalently,

$$p_j(\theta_1, \dots, \theta_k) = p_j(\mu_1, \dots, \mu_k), \quad j = 1, \dots, k.$$

The definition of the power sum $p_j(\dots)$ is given in (4). By Newton's identities (5), we have

$$e_j(\theta_1, \dots, \theta_k) = e_j(\mu_1, \dots, \mu_k), \quad j = 1, \dots, k.$$

Finally, the relation between elementary symmetric polynomials and roots in (3) implies that $\prod_{j=1}^k (t - \theta_j)$ and $\prod_{j=1}^k (t - \mu_j)$ are the same polynomials. Hence, we obtain the conclusion $\ell(\theta, \mu) = 0$. The other direction trivially holds. This verifies the condition of identifiability for the functions f_1, \dots, f_k .

Remark 1 The computation of the statistic $\sum_{j=1}^k f_l(X_j)$ for each $l \in [k]$ is straightforward, thanks to the formula (11). According to (11), we can write

$$\sum_{j=1}^k f_l(X_j) = \sum_{i=1}^k (-1)^{k-i} \frac{e_{k-i}(\mu_1, \dots, \mu_{l-1}, \mu_{l+1}, \dots, \mu_k)}{\prod_{i \in [k] \setminus \{l\}} (\mu_l - \mu_i)} \frac{\sum_{j=1}^k X_j^i}{i}.$$

In other words, $\sum_{j=1}^k f_l(X_j)$ is a linear combination of empirical moments $\{\sum_{j=1}^k X_j^i : i \in [k]\}$. To compute the elementary symmetric polynomial $e_{k-i}(\mu_1, \dots, \mu_{l-1}, \mu_{l+1}, \dots, \mu_k)$ in the coefficient, one can recursively apply Newton's identities (5). The overall complexity of computing $\sum_{j=1}^k f_l(X_j)$ requires $O(k^2)$ products.

We propose the testing statistic

$$T = n \sum_{l=1}^k \left(\sum_{j=1}^k f_l(X_j) - \sum_{j=1}^k f_l(\mu_j) \right)^2. \quad (12)$$

When the value of T is large, the equations (9) are unlikely to hold. Thus, the null hypothesis will be rejected when T exceeds some threshold. The asymptotic distribution of T can be derived under the null hypothesis.

Condition A 1 Assume μ_1, \dots, μ_k are k different numbers that do not vary with n .

Some possible extensions beyond Condition A will be discussed in Section 6.

Theorem 2 Under Condition A, $T \rightsquigarrow \chi_k^2$ as $n \rightarrow \infty$ under the null hypothesis.

For a chi-squared random variable χ_k^2 , define a number $\chi_k^2(\alpha)$ such that

$$\mathbb{P}(\chi_k^2 \leq \chi_k^2(\alpha)) = 1 - \alpha.$$

Then, a testing function is

$$\phi_\alpha = \mathbb{I}\{T > \chi_k^2(\alpha)\}.$$

By Theorem 2, its asymptotic Type-1 error is α . The next result characterizes the regime where the asymptotic power of the test tends to 1. It is a consequence of the fact that the functions f_1, \dots, f_k satisfy the identifiability condition.

Theorem 3 Under Condition A, the following two statements are equivalent

1. $\lim_{n \rightarrow \infty} \sqrt{n} \ell(\theta, \mu) = \infty$;
2. $\lim_{n \rightarrow \infty} \mathbb{P}_\theta(T > \chi_k^2(\alpha)) = 1$, for any constant $\alpha \in (0, 1)$,

where the probability \mathbb{P}_θ denotes $N(\theta, n^{-1}I_k)$.

Theorem 3 shows that $\lim_{n \rightarrow \infty} \sqrt{n} \ell(\theta, \mu) = \infty$ is the necessary and sufficient condition for the asymptotic power of the test to be one. For a local alternative such that $\sqrt{n} \ell(\theta, \mu) = O(1)$, the test will have a non-trivial power between 0 and 1. This contiguous regime will be studied in Section 6.

4. Degeneracy of the Problem

In the last section, we construct a chi-squared test under the assumption that $\min_{j \neq l} |\mu_j - \mu_l| > 0$. When $\min_{j \neq l} |\mu_j - \mu_l| = 0$, the identifiability condition of the functions f_1, \dots, f_k defined in (10) does not hold. We need to construct summary statistics based on new functions in this degenerate case.

Assume there is a partition of the set $[k]$. That is, for some $d \leq k$, we have $\cup_{h=1}^d \mathcal{C}_h = [k]$, and for any $g, h \in [d]$ such that $g \neq h$, $\mathcal{C}_g \cap \mathcal{C}_h = \emptyset$. We assume

$$\mu_j = \nu_h, \text{ for all } j \in \mathcal{C}_h.$$

Moreover, we require that $\min_{g \neq h} |\nu_g - \nu_h| > 0$. To motivate the appropriate functions that we will propose, we consider two extreme cases. The first case is when $d = k$. Then, the condition $\min_{j \neq l} |\mu_j - \mu_l| > 0$ still holds, and we can still use the functions f_1, \dots, f_k defined in (10). The second case is when $d = 1$. This implies $\mu_1 = \mu_2 = \dots = \mu_k = \nu_1$. Then, we can use the function

$$g(t) = (t - \nu_1)^2. \quad (13)$$

This leads to an obvious chi-squared statistic $T_g = n \sum_{j=1}^k g(X_j)$.

For a general d , we need to borrow ideas from both extreme cases. We define functions f_1, \dots, f_d that are modifications from (10). Define

$$f_h(t) = \frac{\int \prod_{g \in [d] \setminus \{h\}} (t - \nu_g)}{\prod_{g \in [d] \setminus \{h\}} (\nu_h - \nu_g)}, \quad h = 1, \dots, d. \quad (14)$$

We also need another function to ensure identifiability. Define

$$g(t) = \frac{\prod_{g=1}^d (t - \nu_g)^2}{\sum_{g=1}^d \prod_{h \in [d] \setminus \{g\}} (t - \nu_h)^2}. \quad (15)$$

The function $g(t)$ is well defined when $d \geq 2$. When $d = 1$, we use the definition given by (13). The following proposition shows that the functions f_1, \dots, f_d, g together ensure identifiability via the equations

$$\sum_{j=1}^k f_h(\theta_j) = \sum_{j=1}^k f_h(\mu_j), \quad h = 1, \dots, d, \quad (16)$$

and

$$\sum_{j=1}^k g(\theta_j) = \sum_{j=1}^k g(\mu_j). \quad (17)$$

Proposition 4 *Assume $\min_{g \neq h} |\nu_g - \nu_h| > 0$. We have the following conclusions.*

1. *When $d = 1$, the equation (17) holds if and only if $\ell(\theta, \mu) = 0$.*
2. *When $2 \leq d \leq k - 1$, the equations (16) and (17) hold if and only if $\ell(\theta, \mu) = 0$.*
3. *When $d = k$, the equation (16) holds if and only if $\ell(\theta, \mu) = 0$.*

The first conclusion of the above proposition is obvious. The last conclusion is proved in Section 3. Here we show the second conclusion. Using a similar argument that we used in Section 3, the equations (16) can be written as

$$E(\nu_1, \dots, \nu_d)\Delta = 0,$$

where $\Delta \in \mathbb{R}^d$ is a vector with the h th entry being $\Delta_h = \frac{1}{h} \sum_{j=1}^k \theta_j^h - \frac{1}{h} \sum_{j=1}^k \mu_j^h$. In other words, we have

$$\sum_{j=1}^k \theta_j^h = \sum_{j=1}^k \mu_j^h, \quad \text{for } h = 1, \dots, d.$$

The equation (17) immediately implies that each θ_j only takes value in the set $\{\nu_1, \dots, \nu_d\}$. Therefore, there exists a partition $[k] = \cup_{h=1}^d \mathcal{D}_h$ such that $\mathcal{D}_g \cap \mathcal{D}_h = \emptyset$ for all $g \neq h$, and $\theta_j = \nu_g$ for all $j \in \mathcal{D}_g$. This leads to

$$\sum_{j=1}^k \theta_j^h = \sum_{g=1}^d |\mathcal{D}_g| \nu_g^h.$$

We also have

$$\sum_{j=1}^k \mu_j^h = \sum_{g=1}^d |\mathcal{C}_g| \nu_g^h.$$

Hence, we obtain the equations

$$\sum_{g=1}^d |\mathcal{D}_g| \nu_g^h = \sum_{g=1}^d |\mathcal{C}_g| \nu_g^h, \quad \text{for } h = 0, 1, \dots, d-1.$$

The equation for $h = 0$ holds because $\sum_{g=1}^d |\mathcal{D}_g| = \sum_{g=1}^d |\mathcal{C}_g| = k$. Again, with matrix notation, these equations can be written as $V(\nu_1, \dots, \nu_d)r = 0$, where $V(\nu_1, \dots, \nu_d)$ is the Vandermonde matrix, and $r \in \mathbb{R}^d$ is a vector with its g th entry being $r_g = |\mathcal{D}_g| - |\mathcal{C}_g|$. When $\min_{g \neq h} |\nu_g - \nu_h| > 0$ holds, $V(\nu_1, \dots, \nu_d)$ has full rank, which leads to $|\mathcal{D}_g| = |\mathcal{C}_g|$ for all $g = 1, \dots, d$. Finally, we can conclude that $\ell(\theta, \mu) = 0$. The other direction is obvious.

The above proof actually shows that the function f_d is not needed when $d \leq k-1$. The equation with $h = d$ in (16) is redundant for identifiability. The second conclusion of Proposition 4 would still hold without it. However, we still keep it for the convenience of analyzing the proposed test.

We propose two testing statistics. Define

$$T_f = n \sum_{h=1}^d \frac{1}{|\mathcal{C}_h|} \left(\sum_{j=1}^k f_h(X_j) - \sum_{j=1}^k f_h(\mu_j) \right)^2, \quad (18)$$

and

$$T_g = n \sum_{j=1}^k g(X_j). \quad (19)$$

We present asymptotic distributions of T_f and T_g . Since the case $d = 1$ is trivial, we only present results for $d \geq 2$.




Figure 1: histograms of testing statistics with $\mu = (1, 3, 3, 3, 5, 5)$, $k = 6$, $d = 3$ and $n = 200$.

Condition B 1 Assume μ_1, \dots, μ_k are k numbers that do not vary with n . Moreover, $\mu_j = \nu_h$ for all $j \in \mathcal{C}_h$, $h \in [d]$.

Theorem 5 Under Condition B, $T_f \rightsquigarrow \chi_d^2$, $T_g \rightsquigarrow \chi_k^2$ and $T_g - T_f \rightsquigarrow \chi_{k-d}^2$ as $n \rightarrow \infty$ under the null hypothesis.

Interestingly, Theorem 5 exhibits an analysis-of-variance type of result. The three statistics all exhibit asymptotic chi-square distributions (see Figure 1). The statistic T_g dominates T_f in probability under the null hypothesis. An analogous analysis-of-variance type of result continues to hold under a local alternative (see Theorem 23). We define the testing function as

$$\phi_\alpha = \mathbb{I}\{T_f > \chi_d^2(\alpha)\} \vee \mathbb{I}\{T_g > \chi_k^2(\alpha)\},$$

where we use the notation $a \vee b$ for $\max(a, b)$. Since under the null hypothesis, $T_g \geq T_f$ in probability, the asymptotic Type-1 error is just the probability of the event $\{T_g > \chi_k^2(\alpha)\}$, which tends to α as $n \rightarrow \infty$. In fact, as we will show later in Section 6, the behavior of the testing function mainly depends on the statistic T_g in the contiguous neighborhood of the null hypothesis. The statistic T_f helps to ensure that the testing function has asymptotic power 1 as soon as $\sqrt{n}\ell(\theta, \mu) \rightarrow \infty$. Without T_f , the identifiability property of the test established in Proposition 4 would break down, and the test would lose power outside of the contiguous neighborhood of the null hypothesis. The next theorem rigorously establishes this fact.

Theorem 6 Under Condition B, the following two statements are equivalent

1. $\lim_{n \rightarrow \infty} \sqrt{n}\ell(\theta, \mu) = \infty$;
2. $\lim_{n \rightarrow \infty} \mathbb{P}_\theta(T_f > \chi_d^2(\alpha) \text{ or } T_g > \chi_k^2(\alpha)) = 1$, for any constant $\alpha \in (0, 1)$,

where the probability \mathbb{P}_θ denotes $N(\theta, n^{-1}I_k)$.

5. The Case of Categorical Distribution

Now we are ready to transfer our wisdom from Gaussian distribution to categorical distribution. Consider i.i.d. observations X_1, \dots, X_n from a categorical distribution (p_1, \dots, p_k) . To be specific, for each $i \in [n]$ and $j \in [k]$, $\mathbb{P}(X_i = j) = p_j$. Throughout this section, we use \mathbb{P}_p to denote the probability distribution of X_1, \dots, X_n . We would like to test whether the k numbers p_1, \dots, p_k are identical to some given q_1, \dots, q_k after a permutation of labels. Introduce a distance between the two vectors p and q ,

$$\ell(p, q) = \min_{\pi \in S_k} 2 \sqrt{\sum_{j=1}^k (\sqrt{p_j} - \sqrt{q_{\pi(j)}})^2}. \quad (20)$$

The hypothesis testing problem is

$$H_0 : \ell(p, q) = 0, \quad H_1 : \ell(p, q) > 0.$$

For each $j \in [k]$, define

$$\hat{p}_j = \frac{1}{n} \sum_{i=1}^n \mathbb{I}\{X_i = j\}.$$

Pearson's chi-squared test (Pearson, 1900) is defined as $\chi^2 = n \sum_{j=1}^k \frac{(\hat{p}_j - q_j)^2}{q_j}$, which is asymptotically distributed as χ_{k-1}^2 when $p = q$. However, this test only works when the null hypothesis is simple. Here, our null hypothesis is composite, and there is uncertainty from the underlying permutation of the labels.

Our idea is to borrow the solution for the Gaussian case in Section 3. Intuitively, the vector $(\hat{p}_1, \dots, \hat{p}_k)$ is asymptotically Gaussian after some normalization. However, the normalization step brings extra difficulty for this problem. In the definition of Pearson's chi-squared test, each \hat{p}_j is normalized by $\sqrt{q_j}$ because of the heteroskedasticity and dependence structure of the vector $(\hat{p}_1, \dots, \hat{p}_k)$. This normalization does not work in our setting because the underlying label is not given, and we do not know which $\sqrt{q_j}$ to use. To overcome this issue, we adopt the technique of variance-stabilizing transformation (Anscombe, 1948), and directly work with $\sqrt{\hat{p}_j}$.

This leads to a modification of the definition of the function $f_l(\cdot)$, and the new definition is given by

$$f_l(t) = \frac{\int \prod_{j \in [k] \setminus \{l\}} (\sqrt{t} - \sqrt{q_j})}{\prod_{j \in [k] \setminus \{l\}} (\sqrt{q_l} - \sqrt{q_j})}, \quad l = 1, \dots, k. \quad (21)$$

The testing statistic is

$$T = 4n \sum_{l=1}^k \left(\sum_{j=1}^k f_l(\hat{p}_j) - \sum_{j=1}^k f_l(q_j) \right)^2. \quad (22)$$

Similar to the discussion in Section 3, when the value of T is large, the equations (9) are unlikely to hold. Thus, the null hypothesis will be rejected when T exceeds some threshold. The asymptotic distribution of T can be derived under the null hypothesis.

Condition C 1 Assume q_1, \dots, q_k are k different numbers in $(0, 1)$ that do not vary with n .

Some possible extensions beyond Condition C will be discussed in Section 6.

Theorem 7 Under Condition C, $T \rightsquigarrow \chi_{k-1}^2$ as $n \rightarrow \infty$ under the null hypothesis.

For a chi-squared random variable χ_{k-1}^2 , define a number $\chi_{k-1}^2(\alpha)$ such that

$$\mathbb{P}(\chi_{k-1}^2 \leq \chi_{k-1}^2(\alpha)) = 1 - \alpha.$$

Then, a testing function is

$$\phi_\alpha = \mathbb{I}\{T > \chi_{k-1}^2(\alpha)\}.$$

By Theorem 7, its asymptotic Type-1 error is α . The next result characterizes the regime where the asymptotic power of the test tends to 1. It is a consequence of the fact that the functions f_1, \dots, f_k satisfy the identifiability condition, though with slightly different definitions.

Theorem 8 Under Condition C, the following two statements are equivalent

1. $\lim_{n \rightarrow \infty} \sqrt{n} \ell(p, q) = \infty$;
2. $\lim_{n \rightarrow \infty} \mathbb{P}_p(T > \chi_{k-1}^2(\alpha)) = 1$, for any constant $\alpha \in (0, 1)$,

where the probability \mathbb{P}_p is defined in the beginning of this section.

Next, we study the degenerate case where the k numbers q_1, \dots, q_k only take d values. There is a partition $[k] = \cup_{h=1}^d \mathcal{C}_h$ such that for any $g \neq h$, $\mathcal{C}_g \cap \mathcal{C}_h = \emptyset$. We assume the following condition.

Condition D 1 Assume q_1, \dots, q_k are k numbers in $(0, 1)$ that do not vary with n . Moreover, $q_j = r_h$ for all $j \in \mathcal{C}_h$, $h \in [d]$.

The approach we take is similar to that in Section 4, assisted with the technique of variance-stabilizing transformation. Define

$$f_h(t) = \frac{\int \prod_{g \in [d] \setminus \{h\}} (\sqrt{t} - \sqrt{r_g})}{\prod_{g \in [d] \setminus \{h\}} (\sqrt{r_h} - \sqrt{r_g})}, \quad h = 1, \dots, d, \quad (23)$$

and

$$g(t) = \frac{\prod_{g=1}^d (\sqrt{t} - \sqrt{r_g})^2}{\sum_{g=1}^d \prod_{h \in [d] \setminus \{g\}} (\sqrt{t} - \sqrt{r_h})^2}. \quad (24)$$

Then, define the testing statistics

$$T_f = 4n \sum_{h=1}^d \frac{1}{|\mathcal{C}_h|} \left(\sum_{j=1}^k f_h(\hat{p}_j) - \sum_{j=1}^k f_h(q_j) \right)^2, \quad (25)$$

and

$$T_g = 4n \sum_{j=1}^k g(\hat{p}_j). \quad (26)$$

The properties of T_f and T_g are given by the following theorem. Again, the case $d = 1$ is trivial, and we only present results for $d \geq 2$.

Theorem 9 *Under Condition D, $T_g \rightsquigarrow \chi_{k-1}^2$, $T_f \rightsquigarrow \chi_{d-1}^2$ and $T_g - T_f \rightsquigarrow \chi_{k-d}^2$ as $n \rightarrow \infty$ under the null hypothesis.*

We define the testing function

$$\phi_\alpha = \mathbb{I}\{T_f > \chi_{d-1}^2(\alpha)\} \vee \mathbb{I}\{T_g > \chi_{k-1}^2(\alpha)\}.$$

By Theorem 9, the Type-1 error of this test converges to α . Though T_f is dominated by T_g under the null hypothesis, both are needed to ensure the power goes to 1 under the alternative.

Theorem 10 *Under Condition D, the following two statements are equivalent*

1. $\lim_{n \rightarrow \infty} \sqrt{n} \ell(p, q) = \infty$;
2. $\lim_{n \rightarrow \infty} \mathbb{P}_p(T_f > \chi_{d-1}^2(\alpha) \text{ or } T_g > \chi_{k-1}^2(\alpha)) = 1$, for any constant $\alpha \in (0, 1)$,

where the probability \mathbb{P}_p is defined in the beginning of this section.

6. Optimality of the Test

In this section, we study the optimality issue of the testing problem from a decision-theoretic perspective. The goal is to understand the fundamental limit of the problem and establish optimality results of the proposed testing procedures. We propose to study the setting where a null hypothesis is tested against a local alternative. This leads to a nontrivial power function and a precise asymptotic characterization of the minimax risk of the test. Depending on whether the data generating process is Gaussian or categorical, and whether the null hypothesis is degenerate or not, the optimality of the test will be studied in four different cases.

6.1 Gaussian Distribution: Non-Degenerate Case

We first consider the non-degenerate situation. That is, we assume that μ_1, \dots, μ_k are k different numbers. In Section 3, we impose the assumption that the k numbers μ_1, \dots, μ_k do not depend on n . This assumption can be made significantly weaker. For two indices j and l that are not equal, define

$$\eta_{jl} = \frac{1}{\mu_j - \mu_l} \prod_{h \in [k] \setminus \{j, l\}} \frac{\mu_l - \mu_h}{\mu_j - \mu_h}.$$

It characterizes the relative difference between μ_j and μ_l in the background of the set $\{\mu_1, \dots, \mu_k\}$.

Condition M1 1 *Assume $\lim_{n \rightarrow \infty} \max_{j \neq l} \frac{|\eta_{jl}|}{\sqrt{n}} = 0$.*

To understand Condition M1, we can interpret $|\eta_{jl}| + |\eta_{lj}|$ as approximately the inverse distance between μ_j and μ_l . Therefore, we allow the possibility that $|\mu_j - \mu_l|$ converges to 0, but not as fast as $n^{-1/2}$. Otherwise, the data cannot tell the difference between $\mu_j \neq \mu_l$

and $\mu_j = \mu_l$, which is equivalent to the degenerate case. Recall that the number k is assumed to be a constant that does not vary with n throughout the paper.

Consider the testing problem

$$H_0 : \theta \in \Theta_0 = \{\theta : \ell(\theta, \mu) = 0\}, \quad H_1 : \theta \in \Theta_\delta = \left\{ \theta : \ell(\theta, \mu) = \frac{\delta}{\sqrt{n}} \right\}. \quad (27)$$

That is, we test the null hypothesis against its contiguous alternative. The choice of H_1 ensures a non-trivial asymptotic power. We measure the testing error via the minimax risk function

$$R_n(k, \delta) = \inf_{0 \leq \phi \leq 1} \left\{ \sup_{\theta \in \Theta_0} \mathbb{P}_\theta \phi + \sup_{\theta \in \Theta_\delta} \mathbb{P}_\theta (1 - \phi) \right\}.$$

The probability symbol \mathbb{P}_θ stands for $N(\theta, n^{-1}I_k)$. Throughout the paper, we assume k and δ are fixed constants independent of n .

We first present the lower bound.

Theorem 11 *Under Condition M1, for sufficiently large n , we have*

$$R_n(k, \delta) \geq \inf_{t > 0} \left(\mathbb{P}(\chi_k^2 > t) + \mathbb{P}(\chi_{k, \delta^2}^2 \leq t) \right).$$

Theorem 11 gives the benchmark of the problem. Using the proposed testing statistic T defined in (12), we can achieve this benchmark.

Theorem 12 *Consider the testing procedure $\phi = \mathbb{I}\{T > t^*\}$, where T is defined in (12), and*

$$t^* = \operatorname{argmin}_{t > 0} \left(\mathbb{P}(\chi_k^2 > t) + \mathbb{P}(\chi_{k, \delta^2}^2 \leq t) \right).$$

Under Condition M1, we have

$$\sup_{\theta \in \Theta_0} \mathbb{P}_\theta \phi + \sup_{\theta \in \Theta_\delta} \mathbb{P}_\theta (1 - \phi) \leq (1 + o(1)) \inf_{t > 0} \left(\mathbb{P}(\chi_k^2 > t) + \mathbb{P}(\chi_{k, \delta^2}^2 \leq t) \right),$$

as $n \rightarrow \infty$.

Theorem 12 characterizes both Type-1 and Type-2 error of the test $\phi = \mathbb{I}\{T > t^*\}$. The conclusion holds for any local alternative with $\delta \in (0, \infty)$. It complements the result of Theorem 3. Combining Theorem 11 and Theorem 12, we conclude that the minimax testing error has the following asymptotic formula

$$R_n(k, \delta) = (1 + o(1)) \inf_{t > 0} \left(\mathbb{P}(\chi_k^2 > t) + \mathbb{P}(\chi_{k, \delta^2}^2 \leq t) \right),$$

and this error can be achieved by the test $\phi = \mathbb{I}\{T > t^*\}$ with some carefully chosen t^* only depending on k and δ .

6.2 Gaussian Distribution: Degenerate Case

Now we consider situations of degeneracy. In Section 4, it is assumed that μ_1, \dots, μ_k only take d different values. This assumption can be relaxed. Here, we assume the k numbers μ_1, \dots, μ_k can be approximately clustered into d groups. Given d different numbers ν_1, \dots, ν_d , for any pair $g \neq h$, define

$$\bar{\eta}_{gh} = \frac{1}{\nu_g - \nu_h} \prod_{l \in [k] \setminus \{g, h\}} \frac{\nu_h - \nu_l}{\nu_g - \nu_l}. \quad (28)$$

Condition M2 1 Assume $\lim_{n \rightarrow \infty} \max_{g \neq h} \frac{|\bar{\eta}_{gh}|}{\sqrt{n}} = 0$ and there is a partition $\mathcal{C}_1, \dots, \mathcal{C}_d$ of $[k]$, such that $\limsup_{n \rightarrow \infty} \max_{1 \leq g \leq d} \max_{j \in \mathcal{C}_g} \sqrt{n} |\mu_j - \nu_g| = 0$.

This condition says that μ_1, \dots, μ_k can be approximately clustered into d groups. The within-group distance is of a smaller order than $n^{-1/2}$, and the between-group distance is of a larger order than $n^{-1/2}$.

Consider the same local testing problem (27). The lower bound of the degenerate setting is given by the following theorem.

Theorem 13 Under Condition M2, $n \rightarrow \infty$, we have

$$R_n(k, \delta) \geq (1 + o(1)) \inf_{t > 0} \left(\mathbb{P}(\chi_k^2 > t) + \mathbb{P}(\chi_{k, \delta^2}^2 \leq t) \right).$$

This lower bound can be achieved asymptotically using the testing statistics T_f and T_g defined in (18) and (19).

Theorem 14 Consider the testing procedure $\phi = \mathbb{I}\{T_f > t^*\} \vee \mathbb{I}\{T_g > t^*\}$, where T_f and T_g are defined in (18) and (19), and

$$t^* = \operatorname{argmin}_{t > 0} \left(\mathbb{P}(\chi_k^2 > t) + \mathbb{P}(\chi_{k, \delta^2}^2 \leq t) \right).$$

Under Condition M2, we have

$$\sup_{\theta \in \Theta_0} \mathbb{P}_\theta \phi + \sup_{\theta \in \Theta_\delta} \mathbb{P}_\theta (1 - \phi) \leq (1 + o(1)) \inf_{t > 0} \left(\mathbb{P}(\chi_k^2 > t) + \mathbb{P}(\chi_{k, \delta^2}^2 \leq t) \right),$$

as $n \rightarrow \infty$.

Theorem 14 shows that the test $\phi = \mathbb{I}\{T_f > t^*\} \vee \mathbb{I}\{T_g > t^*\}$ achieves the optimal error asymptotically under a local alternative. As we will show in Theorem 23, $T_g \geq T_f$ in probability under a local alternative that $\sqrt{n}\ell(\theta, \mu) = \delta \in (0, \infty)$. Therefore, the test $\phi = \mathbb{I}\{T_f > t^*\} \vee \mathbb{I}\{T_g > t^*\}$ is asymptotically equivalent to $\mathbb{I}\{T_g > t^*\}$, and the latter only uses T_g . Though the role of the statistic T_f is negligible for a local alternative, we have already shown in Theorem 6 that as soon as $\sqrt{n}\ell(\theta, \mu) \rightarrow \infty$, the effect of using T_f starts to kick in and it is necessary to use both T_f and T_g for the asymptotic power to approach one.

6.3 Categorical Distribution: Non-Degenerate Case

We study the fundamental limit of testing for the categorical distribution. In Section 5, we assume q_1, \dots, q_k are k different numbers that do not depend on n . In this section, we consider a condition that is significantly weaker. Define

$$\zeta_{jl} = \frac{1}{\sqrt{q_j} - \sqrt{q_l}} \prod_{h \in [k] \setminus \{j, l\}} \frac{\sqrt{q_l} - \sqrt{q_h}}{\sqrt{q_j} - \sqrt{q_h}}.$$

Similar to the definition of η_{jl} , ζ_{jl} characterizes the relative difference between $\sqrt{q_j}$ and $\sqrt{q_l}$ in the background of the set $\{\sqrt{q_1}, \dots, \sqrt{q_k}\}$.

Condition M3 1 Assume $\lim_{n \rightarrow \infty} \max_{j \neq l} \frac{|\zeta_{jl}|}{\sqrt{n}} = 0$ and $\min_{1 \leq j \leq k} nq_j(1 - q_j) \rightarrow \infty$.

Compared with Condition M1, the extra requirement $\min_{1 \leq j \leq k} nq_j(1 - q_j) \rightarrow \infty$ in Condition M3 ensures that each q_j is bounded away from 0 and 1 with a gap at least of order n^{-1} . If this extra requirement does not hold, q_j would be asymptotically equivalent to 0 or 1, which results in a degenerate variance.

Consider the testing problem

$$H_0 : p \in \mathcal{P}_0 = \{p : \ell(p, q) = 0\}, \quad H_1 : p \in \mathcal{P}_\delta = \left\{ p : \ell(p, q) = \frac{\delta}{\sqrt{n}} \right\}. \quad (29)$$

Recall that the distance $\ell(\cdot, \cdot)$ is defined in (20).

We present the lower bound.

Theorem 15 Under Condition M3, as $n \rightarrow \infty$, we have

$$R_n(k, \delta) \geq (1 + o(1)) \inf_{t > 0} \left(\mathbb{P}(\chi_{k-1}^2 > t) + \inf_{\{\delta_1, \delta_2: \delta_1^2 + \delta_2^2 = \delta^2\}} \mathbb{P}(\chi_{k-1, \delta_1}^2 + \delta_2^2 \leq t) \right).$$

Theorem 15 gives the benchmark of the problem. Using the testing statistic T defined in (22), we can achieve this benchmark.

Theorem 16 Consider the testing procedure $\phi = \mathbb{I}\{T > t^*\}$, where T is defined in (22), and

$$t^* = \operatorname{argmin}_{t > 0} \left(\mathbb{P}(\chi_{k-1}^2 > t) + \sup_{\{\delta_1, \delta_2: \delta_1^2 + \delta_2^2 = \delta^2\}} \mathbb{P}(\chi_{k-1, \delta_1}^2 + \delta_2^2 \leq t) \right).$$

Under Condition M3, we have

$$\sup_{\theta \in \mathcal{P}_0} \mathbb{P}_\theta \phi + \sup_{\theta \in \mathcal{P}_\delta} \mathbb{P}_\theta(1 - \phi) \leq (1 + o(1)) \inf_{t > 0} \left(\mathbb{P}(\chi_{k-1}^2 > t) + \sup_{\{\delta_1, \delta_2: \delta_1^2 + \delta_2^2 = \delta^2\}} \mathbb{P}(\chi_{k-1, \delta_1}^2 + \delta_2^2 \leq t) \right),$$

as $n \rightarrow \infty$.

The upper bound given by Theorem 16 does not exactly match the lower bound given by Theorem 15. The difference lies in the Type-2 error. For the lower bound, we get $\inf_{\{\delta_1, \delta_2: \delta_1^2 + \delta_2^2 = \delta^2\}} \mathbb{P}(\chi_{k-1, \delta_1}^2 + \delta_2^2 \leq t)$, while for the upper bound, it is $\sup_{\{\delta_1, \delta_2: \delta_1^2 + \delta_2^2 = \delta^2\}} \mathbb{P}(\chi_{k-1, \delta_1}^2 + \delta_2^2 \leq t)$. These two quantities are close, because for any δ_1 and δ_2 that satisfy $\delta_1^2 + \delta_2^2 = \delta^2$, the expectation of $\chi_{k-1, \delta_1}^2 + \delta_2^2$ is always $k - 1 + \delta^2$. Therefore, the test using the statistic T is nearly optimal.

6.4 Categorical Distribution: Degenerate Case

Finally, we study the categorical distribution with the presence of degeneracy. In Section 5, we consider the situation where q_1, \dots, q_k take d different values. Here, we propose a much weaker condition. Given d different numbers $r_1, \dots, r_d \in (0, 1)$, for any pair $g \neq h$, define

$$\bar{\zeta}_{gh} = \frac{1}{\sqrt{r_g} - \sqrt{r_h}} \prod_{l \in [k] \setminus \{g, h\}} \frac{\sqrt{r_h} - \sqrt{r_l}}{\sqrt{r_g} - \sqrt{r_l}}.$$

Condition M4 1 Assume $\lim_{n \rightarrow \infty} \max_{j \neq l} \frac{|\bar{\zeta}_{jl}|}{\sqrt{n}} = 0$, $\min_{1 \leq j \leq k} nq_j(1 - q_j) \rightarrow \infty$, and there is a partition $\mathcal{C}_1, \dots, \mathcal{C}_d$ of $[k]$, such that $\limsup_{n \rightarrow \infty} \max_{1 \leq g \leq d} \max_{j \in \mathcal{C}_g} \sqrt{n} |\sqrt{q_j} - \sqrt{r_g}| = 0$.

Condition M4 has the same interpretation as Condition M2. The extra requirement $\min_{1 \leq j \leq k} nq_j(1 - q_j) \rightarrow \infty$ is also needed in Condition M3 to prevent the variance from being degenerate.

The lower bound of the local testing problem (29) is given by the next theorem.

Theorem 17 Under Condition M4, as $n \rightarrow \infty$, we have

$$R_n(k, \delta) \geq (1 + o(1)) \inf_{t > 0} \left(\mathbb{P}(\chi_{k-1}^2 > t) + \inf_{\{\delta_1, \delta_2: \delta_1^2 + \delta_2^2 = \delta^2\}} \mathbb{P}(\chi_{k-1, \delta_1}^2 + \delta_2^2 \leq t) \right).$$

For the matching upper bound, we can use the proposed testing statistics T_f and T_g defined in (25) and (26).

Theorem 18 Consider the testing procedure $\phi = \mathbb{I}\{T_f > t^*\} \vee \mathbb{I}\{T_g > t^*\}$, where T_f and T_g are defined in (25) and (26), and

$$t^* = \operatorname{argmin}_{t > 0} \left(\mathbb{P}(\chi_{k-1}^2 > t) + \sup_{\{\delta_1, \delta_2: \delta_1^2 + \delta_2^2 = \delta^2\}} \mathbb{P}(\chi_{k-1, \delta_1}^2 + \delta_2^2 \leq t) \right).$$

Under Condition M4, we have

$$\sup_{\theta \in \mathcal{P}_0} \mathbb{P}_\theta \phi + \sup_{\theta \in \mathcal{P}_\delta} \mathbb{P}_\theta(1 - \phi) \leq (1 + o(1)) \inf_{t > 0} \left(\mathbb{P}(\chi_{k-1}^2 > t) + \sup_{\{\delta_1, \delta_2: \delta_1^2 + \delta_2^2 = \delta^2\}} \mathbb{P}(\chi_{k-1, \delta_1}^2 + \delta_2^2 \leq t) \right),$$

as $n \rightarrow \infty$.

7. Numerical Studies

In this section, we conduct numerical experiments to verify the theoretical properties of the proposed testing procedures. In each of the following scenarios, we compute power functions of α -level tests for $\alpha = 0.05$ with various sample sizes.

Scenario 1. Consider $X \sim N(\theta, n^{-1}I_k)$, and we test the null hypothesis $\ell(\theta, \mu) = 0$ with μ specified as $\mu = (1, 2, 3, 4, 5)$.

Scenario 2. Consider $X \sim N(\theta, n^{-1}I_k)$, and we test the null hypothesis $\ell(\theta, \mu) = 0$ with μ specified as $\mu = (1, 3, 3, 3, 5, 5)$.




Figure 2: Power Curve of Scenario 1

Scenario 3. Consider $X_1, \dots, X_n \sim (p_1, \dots, p_k)$, and we test the null hypothesis $\ell(p, q) = 0$ with q specified as $q = (0.1, 0.2, 0.3, 0.4)$.

Scenario 4. Consider $X_1, \dots, X_n \sim (p_1, \dots, p_k)$, and we test the null hypothesis $\ell(p, q) = 0$ with q specified as $q = (0.1, 0.1, 0.4, 0.4)$.

Scenario 5. Consider $X_1, \dots, X_n \sim (p_1, \dots, p_k)$ and $Y_1, \dots, Y_m \sim (q_1, \dots, q_k)$, and we test the null hypothesis $\ell(p, q) = 0$. We set $p = (0.1, 0.1, 0.4, 0.4)$ and q to be local perturbations of p in a $O(n^{-1/2})$ neighborhood of p .

The numerical results of the five scenarios are summarized in Figures 2-6. The power curves are plotted in the contiguous regimes where $\ell(\theta, \mu) = O(n^{-1/2})$ or $\ell(p, q) = O(n^{-1/2})$. The grey dashed lines correspond to the nominal 0.05 level of the tests.

In Scenario 1, we vary θ in a local $O(n^{-1/2})$ neighborhood of the null hypothesis μ . It is clear that the power function is increasing with respect to $\sqrt{n}\ell(\theta, \mu)$. Moreover, with different sample sizes, the curves match well with each other. This verifies the conclusion of Theorem 3 that the magnitude of $\sqrt{n}\ell(\theta, \mu)$ asymptotically determines the power of the test. We observe in Figure 2 that the power is very close to 1 when $\sqrt{n}\ell(\theta, \mu)$ is greater than 6. The value of the power at $\sqrt{n}\ell(\theta, \mu) = 0$ corresponds to the Type-1 error in the null hypothesis. The actual Type-1 error is slightly greater than the nominal 0.05 level, but the approximations are reasonable for relatively large sample sizes.

Scenario 2 considers a harder null hypothesis with a degenerate $\mu = (1, 3, 3, 3, 5, 5)$. A 0.05-level test studied in Section 4 requires both testing statistics T_f and T_g . Similar to what is observed in Scenario 1, Figure 3 shows that the power approaches 1 at about $\sqrt{n}\ell(\theta, \mu) = 7$, which is predicted by Theorem 6. However, when $\sqrt{n}\ell(\theta, \mu) = 0$, the actual Type-1 errors are consistently larger than the nominal level 0.05, especially when the sample sizes are relatively small.




Figure 3: Power Curve of Scenario 2




Figure 4: Power Curve of Scenario 3




Figure 5: Power Curve of Scenario 4




Figure 6: Power Curve of Scenario 5

Scenario 3 and Scenario 4 consider categorical distributions with a non-degenerate null $q = (0.1, 0.2, 0.3, 0.4)$ and a degenerate null $q = (0.1, 0.1, 0.4, 0.4)$, respectively. Again, Theorem 8 and Theorem 10 are verified by Figure 4 and Figure 5. The actual Type-1 errors are also larger than the nominal one, and are closer to 0.05 with larger sample sizes.

Finally in Scenario 5, we consider experiments of the two-sample test. According to the definitions of the testing statistics in (30) and (31), it is $\sqrt{\frac{2nm}{n+m}}\ell(p, q)$ that determines the power function asymptotically. Figure 6 shows that different power curves well match each other as functions of $\sqrt{\frac{2nm}{n+m}}\ell(p, q)$. As is predicted by Theorem 20, the power is close to 1 at a reasonably large value of $\sqrt{\frac{2nm}{n+m}}\ell(p, q)$.

A common theme in the above numerical results is that the actual Type-1 errors are always larger than the nominal one. We will give an explanation of this phenomenon in Section 9. Roughly speaking, whenever the null exhibits an ambiguous clustering structure, the asymptotic distribution of the testing statistic under the null is a noncentral chi-square distribution. Though a larger sample size helps to make the noncentrality parameter vanish (Figures 2-6), it still results in an estimate of Type-1 error that is too optimistic with a finite sample size. There are potentially two ways to overcome this difficulty. One is to appeal to a second-order correction, and the other is to estimate the noncentrality parameter in the null distribution. We leave this interesting topic as a future project.

8. Two-Sample Test

Consider two categorical distributions (p_1, \dots, p_k) and (q_1, \dots, q_k) . Suppose we observe i.i.d. observations X_1, \dots, X_n from (p_1, \dots, p_k) and i.i.d. observations Y_1, \dots, Y_m from (q_1, \dots, q_k) . We assume that X_1, \dots, X_n are independent of Y_1, \dots, Y_m . The hypothesis testing problem we study in this section is

$$H_0 : \ell(p, q) = 0, \quad H_1 : \ell(p, q) > 0,$$

where the distance $\ell(\cdot, \cdot)$ is defined in (20). The two-sample testing problem is harder than the one-sample version that we have just studied. The major difficulty is that the definitions of the functions (23) and (24) all depend on the values of (p_1, \dots, p_k) and (q_1, \dots, q_k) under the null hypothesis, which is not available anymore in the two-sample scenario.

Our idea is to estimate the unknown (p_1, \dots, p_k) and (q_1, \dots, q_k) from the data, and then construct data-driven versions of (23) and (24).

For each $j \in [k]$, define $\hat{p}_j = \frac{1}{n} \sum_{i=1}^n \mathbb{I}\{X_i = j\}$. Next, we will apply a variable clustering procedure to $(\hat{p}_1, \dots, \hat{p}_k)$. The goal is to find a partition $\underline{\mathcal{C}}_1, \dots, \underline{\mathcal{C}}_d$ of $[k]$ according to

$$j \sim l \quad \text{if} \quad \sqrt{n}|\sqrt{\hat{p}_j} - \sqrt{\hat{p}_l}| \leq \lambda_n.$$

Algorithmically, one can first sort the vector $(\hat{p}_1, \dots, \hat{p}_k)$, and then find the partition sequentially. There exists a permutation $\sigma \in S_k$, such that we can rank the empirical frequencies as $\hat{p}_{\sigma(1)} \leq \hat{p}_{\sigma(2)} \leq \dots \leq \hat{p}_{\sigma(k)}$. Let \hat{j}_1 be the largest j such that $\sqrt{n}|\sqrt{\hat{p}_{\sigma(j)}} - \sqrt{\hat{p}_{\sigma(1)}}| \leq \lambda_n$, where λ_n is some threshold to be specified later. Then, the first cluster is defined as $\underline{\mathcal{C}}_1 = \{\sigma(1), \sigma(2), \dots, \sigma(\hat{j}_1)\}$. Similarly, we can define the second cluster as $\underline{\mathcal{C}}_2 = \{\sigma(\hat{j}_1 +$

1), ..., $\sigma(\hat{j}_2)\}$, where \hat{j}_2 is the largest j such that $\sqrt{n}|\sqrt{\hat{p}_{\sigma(j)}} - \sqrt{\hat{p}_{\sigma(\hat{j}_1)}}| \leq \lambda_n$. We continue this operation until we obtain a partition $\underline{\mathcal{C}}_1, \dots, \underline{\mathcal{C}}_{\underline{d}}$ of $[k]$. Here, \underline{d} is the number of clusters estimated from the data. Now, for each $g \in [\underline{d}]$, we find the center of the cluster by $\sqrt{\underline{r}_g} = \frac{1}{|\underline{\mathcal{C}}_g|} \sum_{j \in \underline{\mathcal{C}}_g} \sqrt{\hat{p}_j}$. With the numbers $\underline{r}_1, \dots, \underline{r}_{\underline{d}}$, we define

$$\underline{f}_h(t) = \frac{\int \prod_{g \in [\underline{d}] \setminus \{h\}} (\sqrt{t} - \sqrt{\underline{r}_g})}{\prod_{g \in [\underline{d}] \setminus \{h\}} (\sqrt{\underline{r}_h} - \sqrt{\underline{r}_g})}, \quad h = 1, \dots, \underline{d},$$

and

$$\underline{g}(t) = \frac{\prod_{g=1}^{\underline{d}} (\sqrt{t} - \sqrt{\underline{r}_g})^2}{\sum_{g=1}^{\underline{d}} \prod_{h \in [\underline{d}] \setminus \{g\}} (\sqrt{t} - \sqrt{\underline{r}_h})^2}.$$

We repeat the above procedure on the observations Y_1, \dots, Y_m . For each $j \in [k]$, define $\hat{q}_j = \frac{1}{m} \sum_{i=1}^m \mathbb{I}\{Y_i = j\}$. Then, apply the same variable clustering procedure on $(\hat{q}_1, \dots, \hat{q}_k)$, and we obtain a partition $\bar{\mathcal{C}}_1, \dots, \bar{\mathcal{C}}_{\bar{d}}$ of $[k]$. For each $g \in [\bar{d}]$, define $\sqrt{\bar{r}_g} = \frac{1}{|\bar{\mathcal{C}}_g|} \sum_{j \in \bar{\mathcal{C}}_g} \sqrt{\hat{q}_j}$. Analogous definitions of \bar{f}_h 's and \bar{g} are given by

$$\bar{f}_h(t) = \frac{\int \prod_{g \in [\bar{d}] \setminus \{h\}} (\sqrt{t} - \sqrt{\bar{r}_g})}{\prod_{g \in [\bar{d}] \setminus \{h\}} (\sqrt{\bar{r}_h} - \sqrt{\bar{r}_g})}, \quad h = 1, \dots, \bar{d},$$

and

$$\bar{g}(t) = \frac{\prod_{g=1}^{\bar{d}} (\sqrt{t} - \sqrt{\bar{r}_g})^2}{\sum_{g=1}^{\bar{d}} \prod_{h \in [\bar{d}] \setminus \{g\}} (\sqrt{t} - \sqrt{\bar{r}_h})^2}.$$

Now we can define testing statistics for this problem:

$$\begin{aligned} T_f &= \frac{2nm}{n+m} \sum_{h=1}^{\underline{d}} \frac{1}{|\underline{\mathcal{C}}_h|} \left(\sum_{j=1}^k \underline{f}_h(\hat{p}_j) - \sum_{j=1}^k \underline{f}_h(\hat{q}_j) \right)^2 \\ &\quad + \frac{2nm}{n+m} \sum_{h=1}^{\bar{d}} \frac{1}{|\bar{\mathcal{C}}_h|} \left(\sum_{j=1}^k \bar{f}_h(\hat{p}_j) - \sum_{j=1}^k \bar{f}_h(\hat{q}_j) \right)^2, \end{aligned} \quad (30)$$

and

$$T_g = \frac{2nm}{n+m} \left(\sum_{j=1}^k \underline{g}(\hat{q}_j) + \sum_{j=1}^k \bar{g}(\hat{p}_j) \right). \quad (31)$$

The asymptotic distributions of the testing statistics under the null distribution are given below.

Condition E 1 *Asume q_1, \dots, q_k are k numbers in $(0, 1)$ that do not vary with n . Moreover, there exists a $d \geq 2$ and a partition $[k] = \cup_{h=1}^d \mathcal{C}_h$, such that $q_j = r_h$ for all $j \in \mathcal{C}_h$, $h \in [d]$.*

Theorem 19 *Assume λ_n is a diverging sequence that satisfies $\lambda_n = o(\sqrt{n})$ and we also assume $\frac{m}{n+m} \rightarrow \beta \in (0, 1)$. Under Condition E, we have*

$$\begin{aligned} T_g &\rightsquigarrow \frac{1}{2}\beta\mathcal{X}_1 + \frac{1}{2}(1-\beta)\mathcal{X}_2 + \mathcal{X}_3, \\ T_f &\rightsquigarrow \mathcal{X}_3, \\ T_g - T_f &\rightsquigarrow \frac{1}{2}\beta\mathcal{X}_1 + \frac{1}{2}(1-\beta)\mathcal{X}_2, \end{aligned}$$

as $n \rightarrow \infty$ under the null hypothesis, where \mathcal{X}_1 , \mathcal{X}_2 and \mathcal{X}_3 are independent random variables distributed as χ_{k-d}^2 , χ_{k-d}^2 and χ_{d-1}^2 , respectively.

Let $\mathcal{X}(\alpha)$ be the number that satisfies $\mathbb{P}\left(\frac{1}{2}\beta\mathcal{X}_1 + \frac{1}{2}(1-\beta)\mathcal{X}_2 + \mathcal{X}_3 > \mathcal{X}(\alpha)\right) = \alpha$. We define the testing function as

$$\phi_\alpha = \mathbb{I}\{T_f > \mathcal{X}(\alpha)\} \vee \mathbb{I}\{T_g > \mathcal{X}(\alpha)\}.$$

Theorem 19 implies that this test has asymptotic Type-1 error α . The next result characterizes the power behavior of the test.

Theorem 20 *Assume λ_n is a diverging sequence that satisfies $\lambda_n = o(\sqrt{n})$ and we also assume $\frac{m}{n+m} \rightarrow \beta \in (0, 1)$. Under Condition E, the following two statements are equivalent*

1. $\lim_{n \rightarrow \infty} \sqrt{n}\ell(p, q) = \infty$;
2. $\lim_{n \rightarrow \infty} \mathbb{P}_{p,q}(T_f > \mathcal{X}(\alpha) \text{ or } T_g > \mathcal{X}(\alpha)) = 1$, for any constant $\alpha \in (0, 1)$,

where the probability $\mathbb{P}_{p,q}$ stands for the joint distribution of $X_1, \dots, X_n, Y_1, \dots, Y_m$.

Theorem 20 assumes Condition E. That is, q_1, \dots, q_k are fixed numbers do that depend on n , and p_1, \dots, p_k are allowed to vary with n . One can also assume an analogous condition for p_1, \dots, p_k as fixed numbers that satisfy Condition E, and allow q_1, \dots, q_k to vary with n .

9. Discussion and Future Directions

The testing procedures that we propose and analyze in this paper critically depend on the structure of null hypothesis. In Section 3, the mean vector $(\mu_1, \dots, \mu_k)^T$ is assumed to consist of k distinct numbers, and the testing statistic is constructed based on the functions f_1, \dots, f_k defined in (10). In Section 4, we assume $(\mu_1, \dots, \mu_k)^T$ consists of k numbers that take values in $\{\nu_1, \dots, \nu_d\}$ for some $d \leq k$. For this degenerate setting, we use the functions f_1, \dots, f_d and g defined in (14) and (15) to construct the testing statistics.

Much weaker assumptions are considered in Section 6. In Section 6.1, we allow $|\mu_j - \mu_l|$ to converge to 0, but require the difference should be of a larger order than $n^{-1/2}$ for every $j \neq l$. This extends the assumption in Section 3 that μ_1, \dots, μ_k are k distinct numbers that do not vary with n . In Section 6.2, we consider the setting where $|\nu_g - \nu_h|$ is of a larger order than $n^{-1/2}$ for every $g \neq h$, and $|\mu_j - \nu_h|$ is of a smaller order than $n^{-1/2}$ for every $j \in \mathcal{C}_h$. This setting extends the assumption used in Section 4. It turns out that the asymptotic distributions of the proposed testing statistics (Theorem 2 and Theorem 5) are still valid under these more general conditions (see Theorem 22 and Theorem 23 in Section 10.1).

However, the conditions in Section 6.1 and Section 6.2 still do not cover all situations. By requiring the within-cluster distance to be of a smaller order than $n^{-1/2}$ and the between-cluster distance to be of a larger order than $n^{-1/2}$, the numbers μ_1, \dots, μ_k enjoy an approximately exact clustering structure, because for each $j \neq l$, we either have $\sqrt{n}|\mu_j - \mu_l| \rightarrow 0$ or $\sqrt{n}|\mu_j - \mu_l| \rightarrow \infty$, depending on whether j and l are in the same cluster or not. A possible situation $\sqrt{n}|\mu_j - \mu_l| \asymp 1$ is excluded.

In this section, we discuss a situation where the clustering structure of the numbers μ_1, \dots, μ_k is ambiguous. Consider a partition $\mathcal{C}_1, \dots, \mathcal{C}_d$ of $[k]$. Define $\nu_h = \frac{1}{|\mathcal{C}_h|} \sum_{j \in \mathcal{C}_h} \mu_j$. Instead of assuming the within-cluster distance is of a smaller order than $n^{-1/2}$, we consider the situation where $n \sum_{h=1}^d \sum_{j \in \mathcal{C}_h} (\mu_j - \nu_h)^2$ is of a constant order. Moreover, we also assume the between-cluster distance $|\nu_g - \nu_h|$ is of a larger order than $n^{-1/2}$ for every $g \neq h$. This is without loss of generality, because if there is some $g \neq h$, such that $|\nu_g - \nu_h| = O(n^{-1/2})$, then \mathcal{C}_g and \mathcal{C}_h can be combined into a single cluster. Recall the definition of $\bar{\eta}_{gh}$ in (28), we formalize this ambiguous clustering structure into the following condition.

Condition M2' 1 For the partition $\mathcal{C}_1, \dots, \mathcal{C}_d$ and clustering centers ν_1, \dots, ν_d defined above, assume $\lim_{n \rightarrow \infty} \max_{g \neq h} \frac{|\bar{\eta}_{gh}|}{\sqrt{n}} = 0$, and $\tau^2 = \lim_{n \rightarrow \infty} n \sum_{h=1}^d \sum_{j \in \mathcal{C}_h} (\mu_j - \nu_h)^2 \in [0, \infty)$.

Note that Condition M2 is a special case of Condition M2' when $\tau^2 = 0$. The next theorem gives the asymptotic distribution of the testing statistics T_f and T_g defined in (18) and (19) under the null hypothesis.

Theorem 21 Assume Condition M2' holds. Then we have $T_g \rightsquigarrow \chi_{k, \tau^2}^2$, $T_f \rightsquigarrow \chi_d^2$ and $T_g - T_f \rightsquigarrow \chi_{k-d, \tau^2}^2$ as $n \rightarrow \infty$ under the null hypothesis $X \sim N(\mu, n^{-1}I_k)$.

It is interesting to see that the asymptotic distribution of T_g is a noncentral chi-squared distribution even under the null hypothesis. The noncentrality parameter τ^2 characterizes the within-cluster distance of μ_1, \dots, μ_k with respect to the partition $\mathcal{C}_1, \dots, \mathcal{C}_d$. Theorem 21 is reduced to Theorem 5 when $\tau^2 = 0$.

Define a number $\chi_{k, \tau^2}^2(\alpha)$ that satisfies $\mathbb{P}(\chi_{k, \tau^2}^2 \leq \chi_{k, \tau^2}^2(\alpha)) = 1 - \alpha$. Then, an α -level testing function is $\phi_\alpha = \mathbb{I}\{T_g > \chi_{k, \tau^2}^2(\alpha)\} \vee \mathbb{I}\{T_f > \chi_d^2(\alpha)\}$. Compared with the null hypothesis where $\tau^2 = 0$, a nonzero τ^2 requires a higher rejection level. This means the test will have less power under a contiguous alternative, compared with the situation where $\tau^2 = 0$. Suppose $\tilde{\tau}^2 = \lim_{n \rightarrow \infty} n \sum_{h=1}^d \sum_{j \in \mathcal{C}_h} (\theta_j - \nu_h)^2 \in (0, \infty)$. Then, one can also show that $T_g \rightsquigarrow \chi_{k, \tilde{\tau}^2}^2$ under the alternative $X \sim N(\theta, n^{-1}I_k)$. Therefore, the test ϕ_α starts to have power when $\tilde{\tau}^2$ exceeds τ^2 . When $\tilde{\tau}^2$ is close to or even smaller than τ^2 , this test will not have any power under the alternative. On the other hand, outside of the contiguous regime where $\sqrt{n}\ell(\theta, \mu) \rightarrow \infty$, we must have $\tilde{\tau}^2 = \infty$, and then the test will have asymptotic power 1.

From what we have just discussed, we can see that the structure of μ_1, \dots, μ_k plays a critical role on the solution of the problem. The discussion also applies to the case of categorical distributions and we can obtain similar conclusions there. Theorem 21 characterizes the asymptotic distribution of the testing statistics when μ_1, \dots, μ_k exhibit an ambiguous clustering structure, right on the edge of degeneracy. This results in a non-trivial behavior

of the power function. Exact characterization of optimality of the testing problem (as what we have done in Section 6) on the edge of degeneracy remains open, and we shall consider this problem as a future project.

Finally, we discussed a list of open problems that can be viewed as natural extensions of the results in the paper.

1. *Growing or infinite support size.* The paper focuses on the case where k is a fixed integer that does not depend on n . The case with a growing k or even $k = \infty$ is of potential importance in many high-dimensional data analysis situations. This requires new techniques because for a probability vector $p = (p_1, \dots, p_k)$ with a growing or an infinite k , many p_j 's have extremely small values.
2. *Testing a parametric family with permutation invariance.* An extension to the null hypothesis (1) is

$$H_0 : p(j) = f_\lambda(\pi(j)) \quad \text{for some } \lambda \in \Lambda \text{ and some } \pi \in S_k.$$

Here, $\{f_\lambda(j)\}$ is a discrete distribution with an unknown parameter $\lambda \in \Lambda$. An example is Poisson(λ). Without the permutation $\pi \in S_k$, the null hypothesis becomes $p = f_\lambda$ for some $\lambda \in \Lambda$, which is a classical goodness-of-fit test of a parametric family Kulperger and Singh (1982); Kocherlakota and Kocherlakota (1986).

3. *Non-asymptotic study of minimax separation.* This paper considers testing procedures that enjoy asymptotic optimality (Section 6). An important theoretical problem is to understand the minimax separation ρ^* for which one can consistently test the null $\ell(p, q) = 0$ against the alternative $\ell(p, q) > \rho$ if and only if $\rho > \rho^*$. With the permutation invariance, the null hypothesis is a non-convex set, which is in contrast to a convex case that was recently studied by Blanchard et al. (2017).
4. *Other group invariance.* Permutation invariance is a special case of group invariance. A more general question is to consider a null hypothesis that is invariant with respect to other group actions. A recent work Perry et al. (2017) considered a group of cyclic shifts. It would be interesting to understand the method of invariance in a general group theoretic framework.

10. Proofs

In this section, we present the proofs of all results in the paper. In Section 10.1, we derive the asymptotic distributions of the proposed testing statistics in various settings. These results are used to derive Theorem 2, Theorem 5, Theorem 7, Theorem 9, Theorem 19 and Theorem 21. Then, in Section 10.2, we analyze the powers of the proposed tests, which include the proofs of Theorem 3, Theorem 6, Theorem 8, Theorem 10 and Theorem 20. Finally, in Section 10.3, we give proofs of all results in Section 6.

10.1 Asymptotic Distribution of the Testing Statistics

We first present and prove four theorems of the proposed testing statistics in various settings.

Theorem 22 *In addition to Condition M1, assume*

$$\lim_{n \rightarrow \infty} \sqrt{n} \ell(\theta, \mu) = \delta \in [0, \infty).$$

Then, as n tends to infinity, $T \rightsquigarrow \chi_{k, \delta^2}^2$.

Proof We first calculate the derivatives of $f_l(t)$. The first derivative is

$$f_l'(t) = \frac{\prod_{j \in [k] \setminus \{l\}} (t - \mu_j)}{\prod_{j \in [k] \setminus \{l\}} (\mu_l - \mu_j)}.$$

Therefore, $f_l'(\mu_l) = 1$. For any $j \neq l$, we give a bound for $\sup_{|t - \mu_j| \leq n^{-1/2} \epsilon} |f_l'(t)|$. The following inequality is useful.

$$|\eta_{lh}| + |\eta_{hl}| = \frac{1}{|\mu_l - \mu_h|} \left(\prod_{j \in [k] \setminus \{l, h\}} \left| \frac{\mu_l - \mu_j}{\mu_h - \mu_j} \right| + \prod_{j \in [k] \setminus \{l, h\}} \left| \frac{\mu_h - \mu_j}{\mu_l - \mu_j} \right| \right) \geq \frac{2}{|\mu_l - \mu_h|}. \quad (32)$$

Note that

$$\begin{aligned} |f_l'(t)| &= \frac{|t - \mu_j|}{|\mu_l - \mu_j|} \left| \prod_{h \in [k] \setminus \{l, j\}} \frac{t - \mu_h}{\mu_l - \mu_h} \right| \\ &= \frac{|t - \mu_j|}{|\mu_l - \mu_j|} \left| \prod_{h \in [k] \setminus \{l, j\}} \left(\frac{t - \mu_j}{\mu_l - \mu_h} + \frac{\mu_j - \mu_h}{\mu_l - \mu_h} \right) \right| \\ &\leq \frac{|t - \mu_j|}{|\mu_l - \mu_j|} 2^{k-2} \left(\left| \prod_{h \in [k] \setminus \{l, j\}} \left(\frac{t - \mu_j}{\mu_l - \mu_h} \right) \right| + \left| \prod_{h \in [k] \setminus \{l, j\}} \left(\frac{\mu_j - \mu_h}{\mu_l - \mu_h} \right) \right| \right) \\ &\leq 2^{k-2} |t - \mu_j|^{k-1} \left(\frac{|\eta_{jl}| + |\eta_{jl}|}{2} \right) \prod_{h \in [k] \setminus \{l, j\}} \left(\frac{|\eta_{lh}| + |\eta_{hl}|}{2} \right) + 2^{k-2} |t - \mu_j| |\eta_{lj}|. \end{aligned}$$

Therefore, we have the bound

$$\kappa_1(\epsilon) = \max_{j \neq l} \sup_{|t - \mu_j| \leq n^{-1/2} \epsilon} |f_l'(t)| \leq 2^{k-2} \left[\max_{j \neq l} \left(\frac{\epsilon |\eta_{jl}|}{\sqrt{n}} \right)^{k-2} + \max_{j \neq l} \left(\frac{\epsilon |\eta_{jl}|}{\sqrt{n}} \right) \right]. \quad (33)$$

The above bound is useful for $k \geq 3$. For $k = 2$, it is easy to see

$$\kappa_1(\epsilon) = \max_{j \neq l} \sup_{|t - \mu_j| \leq n^{-1/2} \epsilon} |f_l'(t)| \leq \max_{j \neq l} \left(\frac{\epsilon |\eta_{jl}|}{\sqrt{n}} \right). \quad (34)$$

The second derivative of $f_l(t)$ is

$$f_l''(t) = \sum_{j \in [k] \setminus \{l\}} \frac{1}{(\mu_l - \mu_j)} \prod_{h \in [k] \setminus \{l, j\}} \frac{t - \mu_h}{\mu_l - \mu_h}.$$

We give a bound for $\sup_{|t-\mu_l| \leq n^{-1/2}\epsilon} |f_l''(t)|$. Similar calculation gives

$$\kappa_2(\epsilon) = \max_{1 \leq l \leq k} \sup_{|t-\mu_l| \leq n^{-1/2}\epsilon} |f_l''(t)| \leq k \max_{j \neq l} |\eta_{jl}| \max_{j \neq l} \left(1 + \frac{\epsilon |\eta_{jl}|}{\sqrt{n}} \right)^{k-2}. \quad (35)$$

Now we are ready to derive the asymptotic distribution of T . We write the observation as $X_j = \theta_j + n^{-1/2}Z_j$, with $Z_j \sim N(0, 1)$ independently. The condition $\lim_{n \rightarrow \infty} \sqrt{n}\ell(\theta, \mu) = \delta$ implies that there is some n_0 , such that for any $n > n_0$, we have

$$n\ell^2(\theta, \mu) \leq C_n \delta^2,$$

where C_n is a sequence that tends to infinity arbitrarily slowly. In particular, we require that C_n satisfies $C_n \rightarrow \infty$ and $\frac{C_n^{3/2} \max_{j \neq l} |\eta_{jl}|}{\sqrt{n}} \rightarrow 0$. The existence of such sequence C_n is guaranteed by the assumption $\frac{\max_{j \neq l} |\eta_{jl}|}{\sqrt{n}} \rightarrow 0$. Thus, there exists a $\pi \in S_k$, possibly depending on n , such that

$$\max_{1 \leq j \leq k} (\theta_j - \mu_{\pi(j)})^2 \leq \frac{C_n \delta^2}{n}.$$

Since k does not depend on n , $\max_{1 \leq j \leq k} Z_j^2 \leq C_n$ with probability that goes to 1. By triangle inequality,

$$\max_{1 \leq j \leq k} |X_j - \mu_{\pi(j)}| \leq \frac{\sqrt{C_n}(1 + \sqrt{\delta^2})}{\sqrt{n}}, \quad (36)$$

with probability that goes to 1. We use Taylor expansion. For j such that $\pi(j) = l$, we have

$$f_l(X_j) - f_l(\mu_{\pi(j)}) = (X_j - \mu_{\pi(j)}) + \frac{1}{2} f_l''(\xi_{jl})(X_j - \mu_{\pi(j)})^2,$$

where we have used the fact that $f_l'(\mu_l) = 1$. For j such that $\pi(j) \neq l$, we have

$$f_l(X_j) - f_l(\mu_{\pi(j)}) = f_l'(\xi_{jl})(X_j - \mu_{\pi(j)}).$$

Therefore,

$$\begin{aligned} & \left| \sum_{j=1}^k f_l(X_j) - \sum_{j=1}^k f_l(\mu_j) - (X_{\pi^{-1}(l)} - \mu_l) \right| \\ & \leq \frac{1}{2} |f_l''(\xi_{\pi^{-1}(l)})| (X_{\pi^{-1}(l)} - \mu_l)^2 + \sum_{j \neq \pi^{-1}(l)} |f_l'(\xi_{jl})| |X_j - \mu_{\pi(j)}|. \end{aligned}$$

The number ξ_{jl} is between X_j and $\mu_{\pi(j)}$, which implies

$$\max_{j,l} |\xi_{jl} - \mu_{\pi(j)}| \leq \max_{1 \leq j \leq k} |X_j - \mu_{\pi(j)}| \leq \frac{\sqrt{C_n}(1 + \sqrt{\delta^2})}{\sqrt{n}}. \quad (37)$$

Using the bounds (33), (34), (35), (36) and (37), we have

$$\begin{aligned}
& \left| \sum_{j=1}^k f_l(X_j) - \sum_{j=1}^k f_l(\mu_j) - (X_{\pi^{-1}(l)} - \mu_l) \right| \\
& \leq \frac{1}{2} \frac{C_n(1 + \sqrt{\delta^2})^2}{n} \kappa_2 \left(\sqrt{C_n}(1 + \sqrt{\delta^2}) \right) \\
& \quad + (k-1) \kappa_1 \left(\sqrt{C_n}(1 + \sqrt{\delta^2}) \right) \frac{\sqrt{C_n}(1 + \sqrt{\delta^2})}{\sqrt{n}}.
\end{aligned} \tag{38}$$

Therefore,

$$\begin{aligned}
& \left| T - n \sum_{l=1}^k (X_{\pi^{-1}(l)} - \mu_l)^2 \right| \\
& \leq n \sum_{l=1}^k \left| \left(\sum_{j=1}^k f_l(X_j) - \sum_{j=1}^k f_l(\mu_j) \right)^2 - (X_{\pi^{-1}(l)} - \mu_l)^2 \right| \\
& \leq 2n \sum_{l=1}^k |X_{\pi^{-1}(l)} - \mu_l| \left| \sum_{j=1}^k f_l(X_j) - \sum_{j=1}^k f_l(\mu_j) - (X_{\pi^{-1}(l)} - \mu_l) \right| \\
& \quad + n \sum_{l=1}^k \left| \sum_{j=1}^k f_l(X_j) - \sum_{j=1}^k f_l(\mu_j) - (X_{\pi^{-1}(l)} - \mu_l) \right|^2 \\
& \leq 2k\sqrt{n}\sqrt{C_n}(1 + \sqrt{\delta^2}) \left| \sum_{j=1}^k f_l(X_j) - \sum_{j=1}^k f_l(\mu_j) - (X_{\pi^{-1}(l)} - \mu_l) \right| \\
& \quad + kn \left| \sum_{j=1}^k f_l(X_j) - \sum_{j=1}^k f_l(\mu_j) - (X_{\pi^{-1}(l)} - \mu_l) \right|^2.
\end{aligned}$$

By (38), the bound for $\left| T - n \sum_{l=1}^k (X_{\pi^{-1}(l)} - \mu_l)^2 \right|$ is of order $\frac{C_n^{3/2} \max_{j \neq l} |\eta_{jl}|}{\sqrt{n}} \rightarrow 0$. Finally, it is easy to see that

$$n \sum_{l=1}^k (X_{\pi^{-1}(l)} - \mu_l)^2 \sim \chi_{k, \delta_n^2}^2,$$

where $\delta_n^2 = n \|\theta - \mu_\pi\|^2 \rightarrow \delta^2$. Therefore, T converges to χ_{k, δ^2}^2 in distribution. \blacksquare

Theorem 23 *In addition to Condition M2, assume*

$$\lim_{n \rightarrow \infty} \sqrt{n} \ell(\theta, \mu) = \delta \in [0, \infty).$$

Then, as n tends to infinity, $T_g \rightsquigarrow \chi_{k, \delta^2}^2$, and $T_g \geq T_f$ in probability. Moreover, if $\delta^2 = 0$, we have $T_g \rightsquigarrow \chi_k^2$, $T_f \rightsquigarrow \chi_d^2$ and $T_g - T_f \rightsquigarrow \chi_{k-d}^2$.

Proof The case $d = 1$ is obvious. We only prove the case $d \geq 2$. Similar to the inequality (32), we have $|\bar{\eta}_{gh}| + |\bar{\eta}_{hg}| \geq \frac{2}{|\nu_g - \nu_h|}$. By Condition M2, we have

$$\frac{\max_{1 \leq g \leq d} \max_{j \in \mathcal{C}_g} |\mu_j - \nu_g|}{\min_{g \neq h} |\nu_g - \nu_h|} = o(1).$$

The observation is $X_j = \theta_j + n^{-1/2}Z_j$ with $Z_j \sim N(0, 1)$ independently. Use the notation $L = \max_{1 \leq g \leq d} \max_{j \in \mathcal{C}_g} \sqrt{n}|\mu_j - \nu_g| = o(1)$. Under the assumption of the theorem, there exists a sequence C_n that satisfies $C_n \rightarrow \infty$, $C_n^2 L \rightarrow 0$ and $\frac{C_n^{3/2} \max_{g \neq h} |\bar{\eta}_{gh}|}{\sqrt{n}} \rightarrow 0$, such that $\max_{1 \leq j \leq k} Z_j^2 \leq C_n$ with probability tending to 1. Similar to the bound (36), the assumption $\lim_{n \rightarrow \infty} \sqrt{n} \ell(\theta, \mu) = \delta < \infty$ implies the existence of $\pi \in S_k$ such that

$$\max_{1 \leq j \leq k} |X_j - \mu_{\pi(j)}| \leq \frac{\sqrt{C_n}(1 + \sqrt{\delta^2})}{\sqrt{n}}.$$

We first study the asymptotic distribution of T_g . Note that

$$\max_{1 \leq g \leq d} \max_{j \in \mathcal{C}_g} |X_{\pi^{-1}(j)} - \nu_g| \leq \frac{\sqrt{C_n}(1 + \sqrt{\delta^2}) + L}{\sqrt{n}}. \quad (39)$$

Together with Condition M2 and the choice of C_n , we can immediately deduce

$$\frac{\max_{1 \leq g \leq d} \max_{j \in \mathcal{C}_g} |X_{\pi^{-1}(j)} - \nu_g|}{\min_{g \neq h} |\nu_g - \nu_h|} \leq \max_{g \neq h} |\bar{\eta}_{gh}| \frac{\sqrt{C_n}(1 + \sqrt{\delta^2}) + L}{\sqrt{n}} = o(1).$$

The function $g(t)$ can be written as

$$\frac{1}{g(t)} = \sum_{g=1}^d \frac{1}{(t - \nu_g)^2}.$$

For each $j \in \mathcal{C}_g$, we have

$$\frac{(X_{\pi^{-1}(j)} - \nu_g)^2}{g(X_{\pi^{-1}(j)})} = 1 + \sum_{h \in [d] \setminus \{g\}} \frac{(X_{\pi^{-1}(j)} - \nu_g)^2}{(X_{\pi^{-1}(j)} - \nu_h)^2}.$$

Thus,

$$\left| \frac{g(X_{\pi^{-1}(j)})}{(X_{\pi^{-1}(j)} - \nu_g)^2} - 1 \right| \leq \frac{\sum_{h \in [d] \setminus \{g\}} \frac{(X_{\pi^{-1}(j)} - \nu_g)^2}{(X_{\pi^{-1}(j)} - \nu_h)^2}}{1 + \sum_{h \in [d] \setminus \{g\}} \frac{(X_{\pi^{-1}(j)} - \nu_g)^2}{(X_{\pi^{-1}(j)} - \nu_h)^2}} \leq \sum_{h \in [d] \setminus \{g\}} \frac{(X_{\pi^{-1}(j)} - \nu_g)^2}{(X_{\pi^{-1}(j)} - \nu_h)^2}, \quad (40)$$

where the bound on the right hand side above can be bounded by

$$\sum_{h \in [d] \setminus \{g\}} \frac{2(X_{\pi^{-1}(j)} - \nu_g)^2}{(\nu_g - \nu_h)^2 - 2(X_{\pi^{-1}(j)} - \nu_h)^2} \leq 4d \max_{g \neq h} |\bar{\eta}_{gh}| \frac{\sqrt{C_n}(1 + \sqrt{\delta^2}) + L}{\sqrt{n}}.$$

Together with (39), we have

$$\begin{aligned}
& |g(X_{\pi^{-1}(j)}) - (X_{\pi^{-1}(j)} - \nu_g)^2| \\
&= (X_{\pi^{-1}(j)} - \nu_g)^2 \left| \frac{g(X_{\pi^{-1}(j)})}{(X_{\pi^{-1}(j)} - \nu_g)^2} - 1 \right| \\
&\leq 4d \max_{g \neq h} |\bar{\eta}_{gh}| \left(\frac{\sqrt{C_n}(1 + \sqrt{\delta^2}) + L}{\sqrt{n}} \right)^3.
\end{aligned}$$

Therefore

$$\begin{aligned}
& \left| T_g - n \sum_{h=1}^d \sum_{j \in \mathcal{C}_h} (X_{\pi^{-1}(j)} - \nu_h)^2 \right| \\
&\leq n \sum_{h=1}^d \sum_{j \in \mathcal{C}_h} |g(X_{\pi^{-1}(j)}) - (X_{\pi^{-1}(j)} - \nu_g)^2| \\
&\leq 4kd \max_{g \neq h} |\bar{\eta}_{gh}| \frac{\left(\sqrt{C_n}(1 + \sqrt{\delta^2}) + L \right)^3}{\sqrt{n}} = o(1).
\end{aligned} \tag{41}$$

For each $j \in \mathcal{C}_h$,

$$\begin{aligned}
& n|(X_{\pi^{-1}(j)} - \nu_h)^2 - (X_{\pi^{-1}(j)} - \mu_j)^2| \\
&\leq n|\nu_h - \mu_j| |X_{\pi^{-1}(j)} - \nu_h + X_{\pi^{-1}(j)} - \mu_j| \\
&\leq L \left(2\sqrt{C_n}(1 + \sqrt{\delta^2}) + L \right) = o(1).
\end{aligned}$$

Thus,

$$\left| T_g - n \sum_{h=1}^d \sum_{j \in \mathcal{C}_h} (X_{\pi^{-1}(j)} - \mu_j)^2 \right| \tag{42}$$

has a bound that tends to 0. Observe that

$$n \sum_{h=1}^d \sum_{j \in \mathcal{C}_h} (X_{\pi^{-1}(j)} - \mu_j)^2 \sim \chi_{k, \delta_n^2}^2,$$

where

$$\delta_n^2 = n \sum_{j=1}^k (\theta_j - \mu_{\pi(j)})^2 \rightsquigarrow \delta^2.$$

Thus, $T_g \rightsquigarrow \chi_{k, \delta^2}^2$.

Next we derive the asymptotic distribution of T_f . Similar to (33), (34) and (35), we also have

$$\kappa_1(\epsilon) = \max_{g \neq h} \sup_{|t - \nu_g| \leq n^{-1/2}\epsilon} |f'_h(t)| \leq 2^{d-2} \left[\max_{g \neq h} \left(\frac{\epsilon |\bar{\eta}_{gh}|}{\sqrt{n}} \right)^{d-2} + \max_{g \neq h} \left(\frac{\epsilon |\bar{\eta}_{gh}|}{\sqrt{n}} \right) \right], \tag{43}$$

for $d \geq 3$,

$$\kappa_1(\epsilon) = \max_{g \neq h} \sup_{|t - \nu_g| \leq n^{-1/2}\epsilon} |f'_h(t)| \leq \max_{g \neq h} \left(\frac{\epsilon |\bar{\eta}_{gh}|}{\sqrt{n}} \right), \quad (44)$$

for $d = 2$, and

$$\kappa_2(\epsilon) = \max_{1 \leq h \leq d} \sup_{|t - \nu_h| \leq n^{-1/2}\epsilon} |f''_h(t)| \leq d \max_{g \neq h} |\bar{\eta}_{gh}| \max_{g \neq h} \left(1 + \frac{\epsilon |\bar{\eta}_{gh}|}{\sqrt{n}} \right)^{d-2}. \quad (45)$$

For any $j \in \mathcal{C}_g$,

$$\begin{aligned} & f_g(X_{\pi^{-1}(j)}) - f_g(\mu_j) \\ &= f_g(X_{\pi^{-1}(j)}) - f_g(\nu_g) + f_g(\nu_g) - f_g(\mu_j) \\ &= f'_g(\nu_g)(X_{\pi^{-1}(j)} - \mu_j) + \frac{1}{2} f''_g(\xi_{jg})(X_{\pi^{-1}(j)} - \nu_g)^2 - \frac{1}{2} f''_g(\bar{\xi}_{jg})(\mu_j - \nu_g)^2. \end{aligned}$$

For any $j \in \mathcal{C}_h$ with any $h \neq g$,

$$f_g(X_{\pi^{-1}(j)}) - f_g(\mu_j) = f'_g(\tilde{\xi}_{jg})(X_{\pi^{-1}(j)} - \mu_j).$$

By the fact that $f'_g(\nu_g) = 1$, we have

$$\begin{aligned} & \left| \sum_{j=1}^k f_g(X_j) - \sum_{j=1}^k f_g(\mu_j) - \sum_{j \in \mathcal{C}_g} (X_{\pi^{-1}(j)} - \mu_j) \right| \\ & \leq \frac{1}{2} \sum_{j \in \mathcal{C}_g} |f''_g(\xi_{jg})| (X_{\pi^{-1}(j)} - \nu_g)^2 + \frac{1}{2} \sum_{j \in \mathcal{C}_g} |f''_g(\bar{\xi}_{jg})| (\mu_j - \nu_g)^2 \\ & \quad \sum_{h \in [d] \setminus \{g\}} \sum_{j \in \mathcal{C}_h} |f'_g(\tilde{\xi}_{jg})| |X_{\pi^{-1}(j)} - \mu_j|. \end{aligned}$$

The number ξ_{jg} is between $X_{\pi^{-1}(j)}$ and ν_g , the number $\bar{\xi}_{jg}$ is between μ_j and ν_g , and the number $\tilde{\xi}_{jg}$ is between $X_{\pi^{-1}(j)}$ and μ_j . Thus,

$$|\xi_{jg} - \nu_g| \leq |X_{\pi^{-1}(j)} - \nu_g| \leq \frac{\sqrt{C_n}(1 + \sqrt{\delta^2}) + L}{\sqrt{n}},$$

$$|\bar{\xi}_{jg} - \nu_g| \leq |\mu_j - \nu_g| \leq \frac{L}{\sqrt{n}},$$

and

$$|\tilde{\xi}_{jg} - \mu_j| \leq |X_{\pi^{-1}(j)} - \mu_j| \leq \frac{\sqrt{C_n}(1 + \sqrt{\delta^2})}{\sqrt{n}}.$$

Using the bounds (43), (44) and (45), we can deduce

$$\begin{aligned} & \left| \sum_{j=1}^k f_g(X_j) - \sum_{j=1}^k f_g(\mu_j) - \sum_{j \in \mathcal{C}_g} (X_{\pi^{-1}(j)} - \mu_j) \right| \\ & \leq k \frac{(\sqrt{C_n}(1 + \sqrt{\delta^2}) + L)^2}{n} \kappa_2 \left(\sqrt{C_n}(1 + \sqrt{\delta^2}) + L \right) \\ & \quad + k \kappa_1 \left(\sqrt{C_n}(1 + \sqrt{\delta^2}) \right) \frac{\sqrt{C_n}(1 + \sqrt{\delta^2})}{\sqrt{n}}. \end{aligned}$$

Similar to the proof of Theorem (32), we can show that

$$\left| T_f - n \sum_{h=1}^d \frac{1}{|\mathcal{C}_h|} \left(\sum_{j \in \mathcal{C}_h} (X_{\pi^{-1}(j)} - \mu_j) \right) \right|^2 \quad (46)$$

has a bound of order $\frac{C_n^{3/2} \max_{j \neq l} |\eta_{jl}|}{\sqrt{n}} \rightarrow 0$. Note that when $\delta^2 = 0$,

$$n \sum_{h=1}^d \frac{1}{|\mathcal{C}_h|} \left(\sum_{j \in \mathcal{C}_h} (X_{\pi^{-1}(j)} - \mu_j) \right)^2 \sim \chi_d^2.$$

Thus, $T_f \rightsquigarrow \chi_d^2$.

Finally, we derive the asymptotic distribution for $T_g - T_f$. The bounds for (42) and (46) imply that

$$\left| T_g - T_f - n \sum_{h=1}^d \sum_{j \in \mathcal{C}_h} (X_{\pi^{-1}(j)} - \mu_j)^2 + n \sum_{h=1}^d \frac{1}{|\mathcal{C}_h|} \left(\sum_{j \in \mathcal{C}_h} (X_{\pi^{-1}(j)} - \mu_j) \right)^2 \right|^2$$

has a bound that tends to zero. Thus, the asymptotic distribution of $T_g - T_f$ is the same as that of

$$\begin{aligned} & n \sum_{h=1}^d \sum_{j \in \mathcal{C}_h} (X_{\pi^{-1}(j)} - \mu_j)^2 - n \sum_{h=1}^d \frac{1}{|\mathcal{C}_h|} \left(\sum_{j \in \mathcal{C}_h} (X_{\pi^{-1}(j)} - \mu_j) \right)^2 \\ &= n \sum_{h=1}^d \sum_{j \in \mathcal{C}_h} \left(X_{\pi^{-1}(j)} - \frac{1}{|\mathcal{C}_h|} \sum_{j \in \mathcal{C}_h} X_{\pi^{-1}(j)} - \left(\mu_j - \frac{1}{|\mathcal{C}_h|} \sum_{j \in \mathcal{C}_h} \mu_j \right) \right)^2, \end{aligned}$$

which is χ_{k-d}^2 when $\delta^2 = 0$. Therefore, $T_g - T_f \rightsquigarrow \chi_{k-d}^2$. Without the condition $\delta^2 = 0$, we can still claim $T_g \geq T_f$ in probability. \blacksquare

Theorem 24 For $\pi = \operatorname{argmin}_{\pi \in S_k} \|\sqrt{p} - \sqrt{q_\pi}\|$, define

$$\delta_1^2 = 4n \sum_{l=1}^k (1 - p_l) \left(\sqrt{p_l} - \sqrt{q_{\pi(l)}} \right)^2, \quad (47)$$

and

$$\delta_2^2 = 4n \sum_{l=1}^k p_l \left(\sqrt{p_l} - \sqrt{q_{\pi(l)}} \right)^2. \quad (48)$$

Assume $\limsup_{n \rightarrow \infty} (\delta_1^2 + \delta_2^2) < \infty$. Then, under Condition M3, $T - \delta_2^2 \rightsquigarrow \chi_{k-1, \delta_1^2}^2$, as n tends to infinity.

Proof The proof is almost the same as that of Theorem 22, and therefore we will omit some overlapping details. Largely speaking, we can replace the t, μ_j, θ_j, X_j by $\sqrt{t}, \sqrt{q_j}, \sqrt{p_j}, \sqrt{\hat{p}_j}$, and most parts in the proof of Theorem 22 will go through. Here are a few different details. We write $\sqrt{\hat{p}_j} = \sqrt{p_j} + n^{-1/2}Z_j/2$, with $Z_j = 2\sqrt{n}(\sqrt{\hat{p}_j} - \sqrt{p_j})$. Condition M3 implies that $\max_{1 \leq j \leq k} Z_j^2 = O_P(1)$. Thus, the inequality (36) in the proof of Theorem 22 can be replaced by $\max_{1 \leq j \leq k} |\sqrt{\hat{p}_j} - \sqrt{q_{\pi(j)}}| \leq \frac{\sqrt{C_n(1+\sqrt{\delta_2^2})}}{\sqrt{n}}$. Then, following the same argument in the proof of Theorem 22, we have

$$\left| T - 4n \sum_{l=1}^k (\sqrt{\hat{p}_l} - \sqrt{q_{\pi(l)}})^2 \right| = o_P(1),$$

and it is sufficient to study the asymptotic distribution of $4n \sum_{l=1}^k (\sqrt{\hat{p}_l} - \sqrt{q_{\pi(l)}})^2$. Let Δ be a vector with the l th entry being $2\sqrt{n}(\sqrt{\hat{p}_l} - \sqrt{q_{\pi(l)}})$. Then, we have $4n \sum_{l=1}^k (\sqrt{\hat{p}_l} - \sqrt{q_{\pi(l)}})^2 = \|Z + \Delta\|^2$. Under Condition M3, $Z \rightsquigarrow N(0, I_k - \sqrt{p}\sqrt{p}^T)$ by Lindeberg's central limit theorem together with an argument of delta's method. Therefore, there exists a random vector W that satisfies $W \rightsquigarrow N(0, I_k)$ and $Z = (I_k - \sqrt{p}\sqrt{p}^T)W$. This gives

$$\begin{aligned} \|Z + \Delta\|^2 &= \|(I_k - \sqrt{p}\sqrt{p}^T)W + (I_k - \sqrt{p}\sqrt{p}^T)\Delta + \sqrt{p}\sqrt{p}^T\Delta\|^2 \\ &= \|(I_k - \sqrt{p}\sqrt{p}^T)W + (I_k - \sqrt{p}\sqrt{p}^T)\Delta\|^2 + \|\sqrt{p}\sqrt{p}^T\Delta\|^2, \end{aligned}$$

where $\|(I_k - \sqrt{p}\sqrt{p}^T)W + (I_k - \sqrt{p}\sqrt{p}^T)\Delta\|^2 \rightsquigarrow \chi_{k-1, \delta_1^2}^2$ and $\|\sqrt{p}\sqrt{p}^T\Delta\|^2 = \delta_2^2$. \blacksquare

Theorem 25 For $\pi = \operatorname{argmin}_{\pi \in S_k} \|\sqrt{p} - \sqrt{q_\pi}\|$, define

$$\delta_1^2 = 4n \sum_{l=1}^k (1 - p_l) \left(\sqrt{p_l} - \sqrt{q_{\pi(l)}} \right)^2,$$

and

$$\delta_2^2 = 4n \sum_{l=1}^k p_l \left(\sqrt{p_l} - \sqrt{q_{\pi(l)}} \right)^2.$$

Assume $\limsup_{n \rightarrow \infty} (\delta_1^2 + \delta_2^2) < \infty$. Then, under Condition M4, $T_g - \delta_2^2 \rightsquigarrow \chi_{k-1, \delta_1^2}^2$, as n tends to infinity. Moreover, $T_g \geq T_f$ in probability. Furthermore, when $\delta_1^2 + \delta_2^2 = 0$, $T_g \rightsquigarrow \chi_{k-1}^2$, $T_f \rightsquigarrow \chi_{d-1}^2$ and $T_g - T_f \rightsquigarrow \chi_{k-d}^2$.

Proof The proof is largely the same as that of Theorem 24. We only need to replace the $t, \mu_j, \theta_j, \nu_h, X_j$ in the proof of Theorem 24 by $\sqrt{t}, \sqrt{q_j}, \sqrt{p_j}, \sqrt{r_h}, \sqrt{\hat{p}_j}$. Then, by the same argument, we have

$$\left| T_g - 4n \sum_{h=1}^d \sum_{j \in C_h} (\sqrt{\hat{p}_{\pi^{-1}(j)}} - \sqrt{q_j})^2 \right| = o_P(1),$$

and

$$\left| T_g - T_f - 4n \sum_{h=1}^d \sum_{j \in \mathcal{C}_h} (\sqrt{\hat{p}_{\pi^{-1}(j)}} - \sqrt{q_j})^2 + 4n \sum_{h=1}^d \frac{1}{|\mathcal{C}_h|} \left(\sum_{j \in \mathcal{C}_h} (\sqrt{\hat{p}_{\pi^{-1}(j)}} - \sqrt{q_j}) \right)^2 \right| = o_P(1).$$

The same argument in the proof of Theorem 24 implies that $T_g - \delta_2^2 \rightsquigarrow \chi_{k-1, \delta_1^2}^2$. The conclusion $T_g \geq T_f$ in probability can be deduced by

$$\begin{aligned} & 4n \sum_{h=1}^d \sum_{j \in \mathcal{C}_h} (\sqrt{\hat{p}_{\pi^{-1}(j)}} - \sqrt{q_j})^2 - 4n \sum_{h=1}^d \frac{1}{|\mathcal{C}_h|} \left(\sum_{j \in \mathcal{C}_h} (\sqrt{\hat{p}_{\pi^{-1}(j)}} - \sqrt{q_j}) \right)^2 \\ &= 4n \sum_{h=1}^d \sum_{j \in \mathcal{C}_h} \left(\sqrt{\hat{p}_{\pi^{-1}(j)}} - \sqrt{q_j} - \frac{1}{|\mathcal{C}_h|} \sum_{j \in \mathcal{C}_h} (\sqrt{\hat{p}_{\pi^{-1}(j)}} - \sqrt{q_j}) \right)^2 \geq 0. \end{aligned}$$

Now we derive the results under the null distribution. Recall the definition of Z_j in the proof of Theorem 24. The asymptotic distributions of T_g , T_f and $T_g - T_f$ are the same of those of

$$\sum_{j=1}^k Z_j^2, \quad \sum_{h=1}^d \frac{1}{|\mathcal{C}_h|} \left(\sum_{j \in \mathcal{C}_h} Z_j \right)^2, \quad \sum_{j=1}^k Z_j^2 - \sum_{h=1}^d \frac{1}{|\mathcal{C}_h|} \left(\sum_{j \in \mathcal{C}_h} Z_j \right)^2,$$

respectively under the null hypothesis. According to the argument in the proof of Theorem 24, $Z = (I_k - \sqrt{q}\sqrt{q}^T)W$ with $W \rightsquigarrow N(0, I_k)$. Therefore, $\sum_{j=1}^k Z_j^2 \rightsquigarrow \chi_{k-1}^2$.

Define a $k \times d$ matrix Q with $Q_{jh} = \frac{1}{\sqrt{|\mathcal{C}_h|}}$ if $j \in \mathcal{C}_h$ and $Q_{jh} = 0$ if $j \notin \mathcal{C}_h$. It is easy to see that QQ^T is a projection matrix and $Q^T Q = I_d$. Define a vector $\gamma \in \mathbb{R}^d$ whose h th entry is $\gamma_h = \sqrt{|\mathcal{C}_h| r_h}$. It is easy to see that γ is a unit vector. Moreover, we have $\sqrt{q} = Q\gamma$. With the new notation, we get

$$\sum_{h=1}^d \frac{1}{|\mathcal{C}_h|} \left(\sum_{j \in \mathcal{C}_h} Z_j \right)^2 = \|Q^T Z\|^2.$$

The covariance of $Q^T Z$ is

$$Q^T (I_k - \sqrt{q}\sqrt{q}^T) Q = I_d - \gamma\gamma^T.$$

Therefore, $\|Q^T Z\|^2 \rightsquigarrow \chi_{d-1}^2$. Finally,

$$\sum_{j=1}^k Z_j^2 - \sum_{h=1}^d \frac{1}{|\mathcal{C}_h|} \left(\sum_{j \in \mathcal{C}_h} Z_j \right)^2 = \|Z\|^2 - \|Q^T Z\|^2 = Z^T (I_k - QQ^T) Z = W^T (I_k - QQ^T) W.$$

Therefore, its asymptotic distribution is χ_{k-d}^2 . ■

The results of Theorem 2, Theorem 5, Theorem 7 and Theorem 9 are special cases of Theorem 22, Theorem 23, Theorem 24 and Theorem 25. Next, we give proofs of Theorem 19 and Theorem 21.

Proof [Proof of Theorem 19] Without loss of generality, we can assume that $p_1 = q_1 \leq p_2 = q_2 \leq \dots \leq p_k = q_k$. This is just to simplify the notation. In general, such a rearrangement can always be done with extra notation of permutations. Then, $\mathcal{C}_g = \{j_g + 1, j_g + 2, \dots, j_{g+1}\}$ for $g \in [d]$. According to the assumption, $\min_{g \neq h} \min_{j \in \mathcal{C}_g} \min_{l \in \mathcal{C}_h} \sqrt{n} |\sqrt{p_j} - \sqrt{p_l}| = o(1)$. Moreover, it is easy to see that $\max_{j \in [k]} \sqrt{n} |\sqrt{\hat{p}_j} - \sqrt{p_j}| = O_P(1)$ and $\max_{j \in [k]} \sqrt{m} |\sqrt{\hat{q}_j} - \sqrt{q_j}| = O_P(1)$. This leads to the conclusion

$$\mathbb{P}(\underline{\mathcal{C}}_g = \bar{\mathcal{C}}_g = \mathcal{C}_g \text{ for all } g \in [d] \text{ and } \underline{d} = \bar{d} = d) \rightarrow 1,$$

under Condition E.

From now on, the analysis is on the event $\{\underline{\mathcal{C}}_g = \bar{\mathcal{C}}_g = \mathcal{C}_g \text{ for all } g \in [d] \text{ and } \underline{d} = \bar{d} = d\}$. Define $\underline{Z}_j = 2\sqrt{n}(\sqrt{\hat{p}_j} - \sqrt{p_j})$ and $\bar{Z}_j = 2\sqrt{m}(\sqrt{\hat{q}_j} - \sqrt{q_j})$ for $j \in [k]$. The definition implies that $\max_{j \in [k]} |\underline{Z}_j| = O_P(1)$ and $\max_{j \in [k]} |\bar{Z}_j| = O_P(1)$. The definitions of \underline{r}_g and \bar{r}_g give

$$2\sqrt{n}(\sqrt{\underline{r}_g} - \sqrt{r_g}) = \frac{1}{|\mathcal{C}_g|} \sum_{j \in \mathcal{C}_g} \underline{Z}_j \quad \text{and} \quad 2\sqrt{m}(\sqrt{\bar{r}_g} - \sqrt{r_g}) = \frac{1}{|\mathcal{C}_g|} \sum_{j \in \mathcal{C}_g} \bar{Z}_j.$$

Given that $p_j = q_j = r_g$ for all $j \in \mathcal{C}_g$, we have $\sqrt{n} |\sqrt{\hat{q}_j} - \sqrt{\underline{r}_g}| = O_P(1)$ and $\sqrt{n} |\sqrt{\hat{p}_j} - \sqrt{\bar{r}_g}| = O_P(1)$ for all $j \in \mathcal{C}_g$. We also have $|\sqrt{\hat{q}_j} - \sqrt{\underline{r}_h}|^{-1} = O_P(1)$ and $|\sqrt{\hat{p}_j} - \sqrt{\bar{r}_h}|^{-1} = O_P(1)$ for all $j \in \mathcal{C}_g$ and $h \neq g$.

We first analyze $\underline{g}(t)$. By its definition,

$$\frac{1}{\underline{g}(t)} = \sum_{h=1}^d \frac{1}{(\sqrt{t} - \sqrt{\underline{r}_h})^2}.$$

Thus, for any $j \in \mathcal{C}_g$,

$$\frac{(\sqrt{\hat{q}_j} - \sqrt{\underline{r}_g})^2}{\underline{g}(\hat{q}_j)} = 1 + \sum_{h \in [d] \setminus \{g\}} \frac{(\sqrt{\hat{q}_j} - \sqrt{\underline{r}_h})^2}{(\sqrt{\hat{q}_j} - \sqrt{\underline{r}_h})^2}.$$

Similar to the argument in (40), we get

$$\left| \frac{\underline{g}(\hat{q}_j)}{(\sqrt{\hat{q}_j} - \sqrt{\underline{r}_g})^2} - 1 \right| \leq \sum_{h \in [d] \setminus \{g\}} \frac{(\sqrt{\hat{q}_j} - \sqrt{\underline{r}_g})^2}{(\sqrt{\hat{q}_j} - \sqrt{\underline{r}_h})^2} = O_P(n^{-1}).$$

With some rearrangements, we get

$$\left| \frac{2nm}{n+m} \sum_{j \in [k]} \underline{g}(\hat{q}_j) - \frac{2nm}{n+m} \sum_{g=1}^d \sum_{j \in \mathcal{C}_g} (\sqrt{\hat{q}_j} - \sqrt{\underline{r}_g})^2 \right| = o_P(1).$$

A similar argument also gives

$$\left| \frac{2nm}{n+m} \sum_{j \in [k]} \bar{g}(\hat{p}_j) - \frac{2nm}{n+m} \sum_{g=1}^d \sum_{j \in \mathcal{C}_g} (\sqrt{\hat{p}_j} - \sqrt{\bar{r}_g})^2 \right| = o_P(1).$$

Therefore, we obtain the following approximation

$$\left| T_g - \frac{2nm}{m+n} \sum_{g=1}^d \sum_{j \in \mathcal{C}_g} \left(\frac{1}{2\sqrt{n}} \underline{Z}_j - \frac{1}{2\sqrt{m}} \frac{1}{|\mathcal{C}_g|} \sum_{j \in \mathcal{C}_g} \bar{Z}_j \right)^2 - \frac{2nm}{m+n} \sum_{g=1}^d \sum_{j \in \mathcal{C}_g} \left(\frac{1}{2\sqrt{m}} \bar{Z}_j - \frac{1}{2\sqrt{n}} \frac{1}{|\mathcal{C}_g|} \sum_{j \in \mathcal{C}_g} \underline{Z}_j \right)^2 \right| = o_P(1).$$

Since

$$\begin{aligned} & \sum_{j \in \mathcal{C}_g} \left(\frac{1}{2\sqrt{n}} \underline{Z}_j - \frac{1}{2\sqrt{m}} \frac{1}{|\mathcal{C}_g|} \sum_{j \in \mathcal{C}_g} \bar{Z}_j \right)^2 \\ &= \sum_{j \in \mathcal{C}_g} \left(\frac{1}{2\sqrt{n}} \underline{Z}_j - \frac{1}{2\sqrt{n}} \frac{1}{|\mathcal{C}_g|} \sum_{j \in \mathcal{C}_g} \underline{Z}_j \right)^2 + |\mathcal{C}_g| \left(\frac{1}{2\sqrt{n}} \frac{1}{|\mathcal{C}_g|} \sum_{j \in \mathcal{C}_g} \underline{Z}_j - \frac{1}{2\sqrt{m}} \frac{1}{|\mathcal{C}_g|} \sum_{j \in \mathcal{C}_g} \bar{Z}_j \right)^2, \end{aligned}$$

and

$$\begin{aligned} & \sum_{j \in \mathcal{C}_g} \left(\frac{1}{2\sqrt{m}} \bar{Z}_j - \frac{1}{2\sqrt{n}} \frac{1}{|\mathcal{C}_g|} \sum_{j \in \mathcal{C}_g} \underline{Z}_j \right)^2 \\ &= \sum_{j \in \mathcal{C}_g} \left(\frac{1}{2\sqrt{m}} \bar{Z}_j - \frac{1}{2\sqrt{m}} \frac{1}{|\mathcal{C}_g|} \sum_{j \in \mathcal{C}_g} \bar{Z}_j \right)^2 + |\mathcal{C}_g| \left(\frac{1}{2\sqrt{n}} \frac{1}{|\mathcal{C}_g|} \sum_{j \in \mathcal{C}_g} \underline{Z}_j - \frac{1}{2\sqrt{m}} \frac{1}{|\mathcal{C}_g|} \sum_{j \in \mathcal{C}_g} \bar{Z}_j \right)^2, \end{aligned}$$

we have

$$\begin{aligned} & \left| T_g - \frac{m}{2(n+m)} \sum_{g=1}^d \sum_{j \in \mathcal{C}_g} \left(\underline{Z}_j - \frac{1}{|\mathcal{C}_g|} \sum_{j \in \mathcal{C}_g} \underline{Z}_j \right)^2 - \frac{n}{2(n+m)} \sum_{g=1}^d \sum_{j \in \mathcal{C}_g} \left(\bar{Z}_j - \frac{1}{|\mathcal{C}_g|} \sum_{j \in \mathcal{C}_g} \bar{Z}_j \right)^2 \right. \\ & \left. - \sum_{g=1}^d |\mathcal{C}_g| \left(\frac{1}{|\mathcal{C}_g|} \sum_{j \in \mathcal{C}_g} \left(\sqrt{\frac{m}{m+n}} \underline{Z}_j - \sqrt{\frac{n}{m+n}} \bar{Z}_j \right) \right)^2 \right| = o_P(1). \end{aligned} \quad (49)$$

Next, we analyze $\underline{f}_h(t)$. By its definition,

$$\frac{d\underline{f}_h(t)}{d\sqrt{t}} = \frac{\prod_{g \in [d] \setminus \{h\}} (\sqrt{t} - \sqrt{r_g})}{\prod_{g \in [d] \setminus \{h\}} (\sqrt{r_h} - \sqrt{r_g})}.$$

Therefore, we have

$$\max_{g \in [d]} \sup_{\sqrt{n}|\sqrt{t} - \sqrt{r_g}| \leq \lambda_n} \left| \frac{d\underline{f}_g(t)}{d\sqrt{t}} - 1 \right| = o_P(1) \quad \text{and} \quad \max_{g \in [d] \setminus \{h\}} \sup_{\sqrt{n}|\sqrt{t} - \sqrt{r_g}| \leq \lambda_n} \left| \frac{d\underline{f}_h(t)}{d\sqrt{t}} \right| = o_P(1).$$

Using Taylor expansion, we get

$$\sum_{j=1}^k f_h(\hat{p}_j) - \sum_{j=1}^k f_h(\hat{q}_j) = \sum_{j \in \mathcal{C}_h} (\sqrt{\hat{p}_j} - \sqrt{\hat{q}_j}) + o_P(1) \sum_{j=1}^k |\sqrt{\hat{p}_j} - \sqrt{\hat{q}_j}|.$$

Then, we have

$$\left| \frac{2nm}{n+m} \sum_{h=1}^d \frac{1}{|\mathcal{C}_h|} \left(\sum_{j=1}^k f_h(\hat{p}_j) - \sum_{j=1}^k f_h(\hat{q}_j) \right)^2 - \frac{2nm}{n+m} \sum_{h=1}^d \frac{1}{|\mathcal{C}_h|} \left(\sum_{j \in \mathcal{C}_h} (\sqrt{\hat{p}_j} - \sqrt{\hat{q}_j}) \right)^2 \right| = o_P(1).$$

The same argument also leads to

$$\left| \frac{2nm}{n+m} \sum_{h=1}^d \frac{1}{|\mathcal{C}_h|} \left(\sum_{j=1}^k \bar{f}_h(\hat{p}_j) - \sum_{j=1}^k \bar{f}_h(\hat{q}_j) \right)^2 - \frac{2nm}{n+m} \sum_{h=1}^d \frac{1}{|\mathcal{C}_h|} \left(\sum_{j \in \mathcal{C}_h} (\sqrt{\hat{p}_j} - \sqrt{\hat{q}_j}) \right)^2 \right| = o_P(1).$$

Hence, we have the following approximation,

$$\left| T_f - \sum_{g=1}^d |\mathcal{C}_g| \left(\frac{1}{|\mathcal{C}_g|} \sum_{j \in \mathcal{C}_g} \left(\sqrt{\frac{m}{m+n}} \underline{Z}_j - \sqrt{\frac{n}{m+n}} \bar{Z}_j \right) \right)^2 \right| = o_P(1). \quad (50)$$

According to the argument in the proof of Theorem 24, $\underline{Z} = (I_k - \sqrt{p}\sqrt{p}^T)\underline{W}$ with $\underline{W} \rightsquigarrow N(0, I_k)$. Similarly, we also have $\bar{Z} = (I_k - \sqrt{q}\sqrt{q}^T)\bar{W}$ with $\bar{W} \rightsquigarrow N(0, I_k)$. Note that \bar{W} is independent of \underline{W} . Recall the definition of the matrix Q and the vector γ in the proof of Theorem 25. Then,

$$\begin{aligned} \sum_{g=1}^d \sum_{j \in \mathcal{C}_g} \left(\underline{Z}_j - \frac{1}{|\mathcal{C}_g|} \sum_{j \in \mathcal{C}_g} \underline{Z}_j \right)^2 &= \underline{Z}^T (I_k - QQ^T) \underline{Z}, \\ \sum_{g=1}^d \sum_{j \in \mathcal{C}_g} \left(\bar{Z}_j - \frac{1}{|\mathcal{C}_g|} \sum_{j \in \mathcal{C}_g} \bar{Z}_j \right)^2 &= \bar{Z}^T (I_k - QQ^T) \bar{Z}, \\ \sum_{g=1}^d |\mathcal{C}_g| \left(\frac{1}{|\mathcal{C}_g|} \sum_{j \in \mathcal{C}_g} \left(\sqrt{\frac{m}{m+n}} \underline{Z}_j - \sqrt{\frac{n}{m+n}} \bar{Z}_j \right) \right)^2 &= \left\| Q^T \left(\sqrt{\frac{m}{m+n}} \underline{Z} - \sqrt{\frac{n}{m+n}} \bar{Z} \right) \right\|^2. \end{aligned}$$

Furthermore, we have

$$\begin{aligned} \underline{Z}^T (I_k - QQ^T) \underline{Z} &= \underline{W}^T (I_k - QQ^T) \underline{W}, \\ \bar{Z}^T (I_k - QQ^T) \bar{Z} &= \bar{W}^T (I_k - QQ^T) \bar{W}, \\ \left\| Q^T \left(\sqrt{\frac{m}{m+n}} \underline{Z} - \sqrt{\frac{n}{m+n}} \bar{Z} \right) \right\|^2 &= \left\| (I_k - \gamma\gamma^T) Q^T \left(\sqrt{\frac{m}{m+n}} \underline{W} - \sqrt{\frac{n}{m+n}} \bar{W} \right) \right\|^2. \end{aligned}$$

Therefore, the three terms above are asymptotically independent, and their asymptotic distributions are χ_{k-d}^2 , χ_{k-d}^2 and χ_{d-1}^2 under the null, respectively. \blacksquare

Proof [Proof of Theorem 21] We will borrow notation and arguments used in the proof of Theorem 23. For example, we keep using the notation $L = \max_{1 \leq g \leq d} \max_{j \in \mathcal{C}_g} \sqrt{n} |\mu_j - \nu_g|$. However, under Condition M2', we have $L = O(1)$ instead of $L = o(1)$. Let C_n be a diverging sequence that satisfies $C_n \rightarrow \infty$ and $\frac{C_n^{3/2} \max_{g \neq h} |\bar{\eta}_{gh}|}{\sqrt{n}} \rightarrow 0$. Then, we can use the same analysis in the proof of Theorem 23 that leads to (41) and (46). Note that the only difference is $L = O(1)$, and it will not affect the conclusions of (41) and (46). We still have

$$\left| T_g - n \sum_{h=1}^d \sum_{j \in \mathcal{C}_h} (X_{\pi^{-1}(j)} - \nu_h)^2 \right| = o_P(1),$$

and

$$\left| T_f - n \sum_{h=1}^d \frac{1}{|\mathcal{C}_h|} \left(\sum_{j \in \mathcal{C}_h} (X_{\pi^{-1}(j)} - \mu_j) \right)^2 \right| = o_P(1).$$

By the fact that

$$\begin{aligned} & n \sum_{h=1}^d \sum_{j \in \mathcal{C}_h} (X_{\pi^{-1}(j)} - \nu_h)^2 - n \sum_{h=1}^d \frac{1}{|\mathcal{C}_h|} \left(\sum_{j \in \mathcal{C}_h} (X_{\pi^{-1}(j)} - \mu_j) \right)^2 \\ &= n \sum_{h=1}^d \sum_{j \in \mathcal{C}_h} (X_{\pi^{-1}(j)} - \nu_h)^2 - n \sum_{h=1}^d |\mathcal{C}_h| \left(\frac{1}{|\mathcal{C}_h|} \sum_{j \in \mathcal{C}_h} (X_{\pi^{-1}(j)} - \nu_h) \right)^2 \\ &= n \sum_{h=1}^d \sum_{j \in \mathcal{C}_h} \left(X_{\pi^{-1}(j)} - \frac{1}{|\mathcal{C}_h|} \sum_{j \in \mathcal{C}_h} X_{\pi^{-1}(j)} \right)^2, \end{aligned}$$

we also have

$$\left| T_g - T_f - n \sum_{h=1}^d \sum_{j \in \mathcal{C}_h} \left(X_{\pi^{-1}(j)} - \frac{1}{|\mathcal{C}_h|} \sum_{j \in \mathcal{C}_h} X_{\pi^{-1}(j)} \right)^2 \right| = o_P(1).$$

Therefore, under the null hypothesis $X \sim N(\mu, n^{-1}I_k)$, we have $T_g \rightsquigarrow \chi_{k, \tau^2}^2$, $T_f \rightsquigarrow \chi_d^2$ and $T_g - T_f \rightsquigarrow \chi_{k-d, \tau^2}^2$. \blacksquare

10.2 Power Analysis

In this section, we give proofs of Theorem 3, Theorem 6, Theorem 8, Theorem 10 and Theorem 20.

Proof [Proof of Theorem 3] We first assume $n\ell(\theta, \mu)^2 \rightarrow \infty$ and derive $T \rightarrow \infty$ in probability. Note that for each $\pi \in S_k$,

$$n \sum_{j=1}^k (\theta_j - \mu_{\pi(j)})^2 \leq 2n \sum_{j=1}^k (X_j - \theta_j)^2 + 2n \sum_{j=1}^k (X_j - \mu_{\pi(j)})^2.$$

Therefore,

$$n\ell(\theta, \mu)^2 \leq 2 \sum_{j=1}^k Z_j^2 + 2n\ell(X, \mu)^2,$$

where $Z_j \sim N(0, 1)$. The fact that $2 \sum_{j=1}^k Z_j^2 = O_P(1)$ and the assumption $n\ell(\theta, \mu)^2 \rightarrow \infty$ implies that $n\ell(X, \mu)^2 \rightarrow \infty$ in probability. Suppose we can show $T = O_P(1)$ implies $n\ell(X, \mu)^2 = O_P(1)$, then $n\ell(X, \mu)^2 \rightarrow \infty$ in probability must implies $T \rightarrow \infty$ in probability.

Now we suppose a bound $T \leq B = O(1)$, and it is sufficient to derive a bound for $n\ell(X, \mu)^2$. For each $j = 1, \dots, k$, we shorthand the power sums $p_j(X_1, \dots, X_k)$ and $p_j(\mu_1, \dots, \mu_k)$ by $p_j(X)$ and $p_j(\mu)$. Similarly, the elementary symmetric polynomials $e_j(X_1, \dots, X_k)$ and $e_j(\mu_1, \dots, \mu_k)$ are shorthanded by $e_j(X)$ and $e_j(\mu)$. Define a vector $\Delta \in \mathbb{R}^k$ with the j th entry being $\Delta_j = \frac{1}{j} \sum_{h=1}^k X_h^j - \frac{1}{j} \sum_{h=1}^k \mu_h^j$. Recall the definition of the matrix $E(\mu_1, \dots, \mu_k)$. Then,

$$T = n\|E(\mu_1, \dots, \mu_k)\Delta\|^2.$$

We use $\lambda_{\max}(\cdot)$ and $\lambda_{\min}(\cdot)$ to denote the largest and the smallest eigenvalues. By the fact that $V(\mu_1, \dots, \mu_k)E(\mu_1, \dots, \mu_k) = I_k$, we have

$$T \geq n\lambda_{\min}(E(\mu_1, \dots, \mu_k)^T E(\mu_1, \dots, \mu_k))\|\Delta\|^2 \geq \frac{n\|\Delta\|^2}{\lambda_{\max}(V(\mu_1, \dots, \mu_k)^T V(\mu_1, \dots, \mu_k))}.$$

The bound $T \leq B$ then leads to

$$\|\Delta\|^2 \leq \frac{\lambda_{\max}(V(\mu_1, \dots, \mu_k)^T V(\mu_1, \dots, \mu_k))B}{n} = O\left(\frac{B}{n}\right). \quad (51)$$

Therefore, $|p_j(X) - p_j(\mu)|^2 = O(B/n)$ for each $j \in [k]$. By Newton's identities, we can deduce $|e_j(X) - e_j(\mu)|^2 = O(B/n)$ for each $j \in [k]$. Define

$$f(t) = \prod_{j=1}^k (t - \mu_j), \quad \hat{f}(t) = \prod_{j=1}^k (t - X_j).$$

The relation between the two polynomials and the elementary symmetric polynomials is given in (3). Using (3), we give a bound for $|f(X_l)|$.

$$|f(X_l)| = |f(X_l) - \hat{f}(X_l)| \leq \sum_{j=0}^k |e_{k-j}(X) - e_{k-j}(\mu)| |X_l|^j.$$

Since $|X_l|^2 \leq p_2(X) \leq p_2(\mu) + |p_2(X) - p_2(\mu)| = O(1)$, we have $|f(X_l)|^2 = O(B/n)$. The following proposition is useful and will be proved in the end.

Proposition 26 *For any μ_1, \dots, μ_k , we have*

$$|f(t)| \geq \min_{1 \leq j \leq k} |t - \mu_j| \prod_{1 \leq j < l \leq k} \left| \frac{\mu_j - \mu_l}{2} \right|.$$

By this inequality, we have

$$\max_{1 \leq l \leq k} \min_{1 \leq j \leq k} (X_l - \mu_j)^2 = O\left(\frac{B}{n}\right). \quad (52)$$

Therefore, there exists a sequence $\sigma(1), \dots, \sigma(k)$ such that

$$\max_{1 \leq j \leq k} (X_j - \mu_{\sigma(j)})^2 = O\left(\frac{B}{n}\right).$$

Since

$$\prod_{j=1}^k |t - \mu_{\sigma(j)}| \leq 2^k \prod_{j=1}^k |t - X_j| + 2^k \prod_{j=1}^k |X_j - \mu_{\sigma(j)}| = 2^k |\hat{f}(t)| + O\left(\left(\frac{B}{n}\right)^{k/2}\right),$$

and

$$|\hat{f}(\mu_l)| = |\hat{f}(\mu_l) - f(\mu_l)| \leq \sum_{j=0}^k |e_{k-j}(X) - e_{k-j}(\mu)| |\mu_l|^j = O\left(\sqrt{\frac{B}{n}}\right),$$

we have

$$\prod_{j=1}^k |\mu_l - \mu_{\sigma(j)}| = O\left(\sqrt{\frac{B}{n}}\right),$$

which holds for every $l = 1, \dots, k$. The fact that μ_1, \dots, μ_k are k different fixed number implies σ must be an element of S_k . Hence, the bound (52) implies $n\ell(X, \mu)^2 = O(B)$, and the proof of one direction is complete.

For the other direction, it is sufficient to show that $n\ell(\theta, \mu) = O(1)$ implies $T = O_P(1)$. This can be shown using the same argument in the proof of Theorem 22. \blacksquare

Proof [Proof of Proposition 26] We first consider the case $k = 2$, where $f(t) = (t - \mu_1)(t - \mu_2)$. Suppose $|t - \mu_1| \leq |t - \mu_2|$, then $|t - \mu_2| \geq \frac{|\mu_1 - \mu_2|}{2}$. Thus, $|f(t)| \geq \frac{|\mu_1 - \mu_2|}{2} \min\{|t - \mu_1|, |t - \mu_2|\}$. The same argument also works when $|t - \mu_1| > |t - \mu_2|$. When $k = 3$,

$$\begin{aligned} |f(t)| &\geq |t - \mu_3| \frac{|\mu_1 - \mu_2|}{2} \min\{|t - \mu_1|, |t - \mu_2|\} \\ &= \frac{|\mu_1 - \mu_2|}{2} \min\{|t - \mu_1||t - \mu_3|, |t - \mu_2||t - \mu_3|\}. \end{aligned}$$

The inequality for $k = 2$ can be used to lower bound both $|t - \mu_1||t - \mu_3|$ and $|t - \mu_2||t - \mu_3|$. This gives the desired result for $k = 3$. A standard mathematical induction argument leads to inequality for all k . \blacksquare

Proof [Proof of Theorem 6] According to the argument that we have used in the proof of Theorem 3, we need to show $T_f = O_P(1)$ and $T_g = O_P(1)$ imply $n\ell(X, \mu)^2 = O_P(1)$ for the proof of the first direction.

Suppose $T_f \leq B_1 = O(1)$ and $T_g \leq B_2 = O(1)$. It is sufficient to derive a bound for $n\ell(X, \mu)^2$. We first derive an inequality for $g(t)$. Since

$$\max_{1 \leq g \leq d} \prod_{h \in [d] \setminus \{g\}} (t - \nu_h)^2 \leq \sum_{g=1}^d \prod_{h \in [d] \setminus \{g\}} (t - \nu_h)^2 \leq d \max_{1 \leq g \leq d} \prod_{h \in [d] \setminus \{g\}} (t - \nu_h)^2,$$

we have

$$\frac{1}{d} \min_{1 \leq g \leq d} (t - \nu_g)^2 \leq g(t) \leq \min_{1 \leq g \leq d} (t - \nu_g)^2. \quad (53)$$

Therefore, $T_g \leq B_2$ implies that $\sum_{j=1}^k \min_{1 \leq g \leq d} (X_j - \nu_g)^2 \leq \frac{dB_2}{n}$. This implies the existence of a sequence $\sigma(1), \dots, \sigma(k)$ such that $\max_{1 \leq j \leq k} (X_j - \nu_{\sigma(j)})^2 \leq \frac{dB_2}{n} = O(B_2/n)$. It further implies $\max_{1 \leq h \leq d} \max_{1 \leq j \leq k} |X_j^h - \nu_{\sigma(j)}^h| = O(\sqrt{B_2/n})$. Define $\hat{\mathcal{C}}_g = \{j \in [k] : \sigma(j) = g\}$ for each $g \in [d]$. Then

$$\sum_{h=1}^{d-1} \left(\sum_{j=1}^k X_j^h - \sum_{g=1}^d |\hat{\mathcal{C}}_g| \nu_g^h \right)^2 = O\left(\frac{B_2}{n}\right).$$

Using the same argument in deriving (51), we can also get the bound

$$\sum_{h=1}^{d-1} \left(\sum_{j=1}^k X_j^h - \sum_{g=1}^d |\mathcal{C}_g| \nu_g^h \right)^2 = O\left(\frac{B_1}{n}\right).$$

The inequalities in the last two displays, together with the equality $\sum_{g=1}^d |\hat{\mathcal{C}}_g| = \sum_{g=1}^d |\mathcal{C}_g|$, give

$$\sum_{h=0}^{d-1} \left(\sum_{g=1}^d |\hat{\mathcal{C}}_g| \nu_g^h - \sum_{g=1}^d |\mathcal{C}_g| \nu_g^h \right)^2 = O\left(\frac{B_1 + B_2}{n}\right).$$

Define a vector $r \in \mathbb{R}^d$, with its g th entry being $|\hat{\mathcal{C}}_g| - |\mathcal{C}_g|$. Then,

$$\sum_{h=0}^{d-1} \left(\sum_{g=1}^d |\hat{\mathcal{C}}_g| \nu_g^h - \sum_{g=1}^d |\mathcal{C}_g| \nu_g^h \right)^2 = \|V(\nu_1, \dots, \nu_d)r\|^2 \geq \lambda_{\min}(V(\nu_1, \dots, \nu_d)^T V(\nu_1, \dots, \nu_d)) \|r\|^2.$$

When ν_1, \dots, ν_d are d different numbers, we have $\lambda_{\min}(V(\nu_1, \dots, \nu_d)^T V(\nu_1, \dots, \nu_d)) > 0$, and thus $\|r\|^2 = O\left(\frac{B_1 + B_2}{n}\right)$. Since $\|r\|^2$ is an integer, we must have $\|r\|^2 = 0$, which gives $|\hat{\mathcal{C}}_g| = |\mathcal{C}_g|$ for any $g \in [d]$. From this we can deduce that $n\ell(X, \mu)^2 = O(B_2)$.

For the other direction, it is sufficient to show that $n\ell(\theta, \mu)^2 = O(1)$ implies $T_f = O_P(1)$ and $T_g = O_P(1)$. This can be shown using the same argument in the proof of Theorem 23. ■

Proof [Proofs of Theorem 8 and Theorem 10] The proofs are the same as those of Theorem 3 and Theorem 6. ■

Proof [Proof of Theorem 20] First of all, we have $\max_{j \in [k]} \sqrt{n} |\sqrt{\hat{q}_j} - \sqrt{q_j}| = O_P(1)$. This gives that $\mathbb{P}(\bar{\mathcal{C}}_g = \mathcal{C}_g \text{ for all } g \in [d] \text{ and } \bar{d} = d) \rightarrow 1$. From now on, the analysis is on the event $\{\bar{\mathcal{C}}_g = \mathcal{C}_g \text{ for all } g \in [d] \text{ and } \bar{d} = d\}$. Since we also have $\max_{j \in [k]} \sqrt{n} |\sqrt{\hat{p}_j} - \sqrt{p_j}| = O_P(1)$, the statement $n\ell(p, q)^2 \rightarrow \infty$ is equivalent to $n\ell(\hat{p}, q)^2 \rightarrow \infty$ in probability. Therefore, we only need to establish the equivalence between $n\ell(\hat{p}, q)^2 \rightarrow \infty$ in probability and the power of the test goes to one.

In the first direction of the proof, we suppose that $T_f \leq B_1 = O_P(1)$ and $T_g \leq B_2 = O_P(1)$, and we will show $n\ell(\hat{p}, q)^2 = O_P(1)$. The bound $T_g \leq B_2 = O_P(1)$ implies that

$$\frac{2nm}{n+m} \sum_{j=1}^k \bar{g}(\hat{p}_j) \leq B_2.$$

By the definition of $\bar{g}(\cdot)$, we have $\bar{g}(t) \geq d^{-1} \min_{g \in [d]} (\sqrt{t} - \sqrt{r_g})^2$. This implies the bound

$$\sum_{j=1}^k \min_{g \in [d]} (\sqrt{\hat{p}_j} - \sqrt{r_g})^2 \leq \frac{dB_2(n+m)}{nm} = O_P(n^{-1}).$$

Since $\max_{g \in [d]} \sqrt{n} |\sqrt{\bar{r}_g} - \sqrt{r_g}| = O_P(1)$, we deduce

$$\sum_{j=1}^k \min_{g \in [d]} (\sqrt{\hat{p}_j} - \sqrt{r_g})^2 = O_P(n^{-1}).$$

Then, there must exist $\sigma(1), \dots, \sigma(k)$ such that $\max_{j \in [k]} |\sqrt{\hat{p}_j} - \sqrt{r_{\sigma(j)}}|^2 = O_P(n^{-1})$. It further implies that $\max_{h \in [d]} \max_{j \in [k]} |(\sqrt{\hat{p}_j})^h - (\sqrt{r_{\sigma(j)}})^h| = O_P(n^{-1/2})$. Define $\hat{\mathcal{C}}_g = \{j \in [k] : \sigma(j) = g\}$ for each $g \in [d]$. Then, we have

$$\sum_{h=1}^{d-1} \left(\sum_{j=1}^k (\sqrt{\hat{p}_j})^h - \sum_{g=1}^d |\hat{\mathcal{C}}_g| (\sqrt{r_g})^h \right)^2 = O_P(n^{-1}). \quad (54)$$

Note that

$$\begin{aligned} T_f &\geq \frac{2nm}{n+m} \sum_{h=1}^d \frac{1}{|\mathcal{C}_h|} \left(\sum_{j=1}^k \bar{f}_h(\hat{p}_j) - \sum_{j=1}^k \bar{f}_h(\hat{q}_j) \right)^2 \\ &\geq \frac{2nm}{d(n+m)} \sum_{h=1}^d \left(\sum_{j=1}^k \bar{f}_h(\hat{p}_j) - \sum_{j=1}^k \bar{f}_h(\hat{q}_j) \right)^2 \\ &= \frac{2nm}{d(n+m)} \|E(\sqrt{\bar{r}_1}, \dots, \sqrt{\bar{r}_d}) \Delta\|^2, \end{aligned}$$

where Δ is a d -dimensional vector with $\Delta_h = \frac{1}{h} \sum_{j=1}^k (\sqrt{\hat{p}_j})^h - \frac{1}{h} \sum_{j=1}^k (\sqrt{\hat{q}_j})^h$. Thus, the bound $T_f \leq B_1 = O_P(1)$ implies that

$$\|E(\sqrt{\bar{r}_1}, \dots, \sqrt{\bar{r}_d}) \Delta\|^2 = O_P(n^{-1}).$$

Since $\lambda_{\min}(E(\sqrt{r_1}, \dots, \sqrt{r_d})^T E(\sqrt{r_1}, \dots, \sqrt{r_d}))$ is a positive constant that is bounded away from 0, and

$$|\lambda_{\min}(E(\sqrt{\hat{r}_1}, \dots, \sqrt{\hat{r}_d})^T E(\sqrt{\hat{r}_1}, \dots, \sqrt{\hat{r}_d})) - \lambda_{\min}(E(\sqrt{r_1}, \dots, \sqrt{r_d})^T E(\sqrt{r_1}, \dots, \sqrt{r_d}))| = o_P(1),$$

we have $\|\Delta\|^2 = O_P(n^{-1})$, which further leads to

$$\sum_{h=1}^{d-1} \left(\sum_{j=1}^k (\sqrt{\hat{p}_j})^h - \sum_{g=1}^d |\mathcal{C}_g| (\sqrt{r_g})^h \right)^2 = O_P(n^{-1}), \quad (55)$$

by using the fact that $\max_{j \in [k]} \sqrt{n} |\sqrt{\hat{q}_j} - \sqrt{q_j}| = O_P(1)$ and Condition E. The two inequalities (54) and (55), together with the fact that $\sum_{g=1}^d |\hat{\mathcal{C}}_g| = \sum_{g=1}^d |\mathcal{C}_g|$, imply

$$\sum_{h=0}^{d-1} \left(\sum_{g=1}^d |\hat{\mathcal{C}}_g| (\sqrt{r_g})^h - \sum_{g=1}^d |\mathcal{C}_g| (\sqrt{r_g})^h \right)^2 = O_P(n^{-1}).$$

The same argument used in the proof of Theorem 6 implies that $|\hat{\mathcal{C}}_g| = |\mathcal{C}_g|$ for all $g \in [d]$. Therefore, together with $\max_{j \in [k]} |\sqrt{\hat{p}_j} - \sqrt{r_{\sigma(j)}}|^2 = O_P(n^{-1})$, we obtain the conclusion $n\ell(\hat{p}, q) = O_P(1)$.

For the other direction, when $n\ell(p, q)^2 = O(1)$, the approximations (49) and (50) in the proofs of Theorem 19 hold with bounds at the order of $O_P(1)$. This leads to $T_f = O_P(1)$ and $T_g = O_P(1)$. \blacksquare

10.3 Minimax Upper and Lower Bounds

In this section, we prove all results in Section 6. We first give proofs for the lower bounds, and then for the upper bounds.

Proof [Proof of Theorem 11] We first observe an inequality $|\eta_{jl}| + |\eta_j| \geq \frac{2}{|\mu_l - \mu_j|}$, which has been derived in the proof of Theorem 22. Thus, Condition M1 implies $\sqrt{n} |\mu_j - \mu_l| \rightarrow \infty$ for any $j \neq l$. Consider the set

$$\bar{\Theta}_\delta = \left\{ \theta : \|\theta - \mu\| = \frac{\delta}{\sqrt{n}} \right\}.$$

For each $\theta \in \bar{\Theta}_\delta$, $|\theta_j - \mu_j|^2 \leq \frac{\delta^2}{n}$, which implies μ_j is the closest element to θ_j in the set $\{\mu_1, \dots, \mu_k\}$. Therefore, $\ell(\theta, \mu) = \|\theta - \mu\| = \delta/\sqrt{n}$, which implies $\bar{\Theta}_\delta \subset \Theta_\delta$. This gives the lower bound

$$R_n(k, \delta) \geq \inf_{0 \leq \phi \leq 1} \left\{ \mathbb{P}_\mu \phi + \sup_{\theta \in \bar{\Theta}_\delta} \mathbb{P}_\theta (1 - \phi) \right\}.$$

Consider the uniform distribution Π on $\bar{\Theta}_\delta$. Then,

$$R_n(k, \delta) \geq \inf_{0 \leq \phi \leq 1} \left\{ \mathbb{P}_\mu \phi + \int \mathbb{P}_\theta (1 - \phi) d\Pi(\theta) \right\}.$$

By Neyman-Pearson lemma, the optimal testing function ϕ is given by

$$\phi = \left\{ \frac{d \int \mathbb{P}_\theta d\Pi(\theta)}{d\mathbb{P}_\mu} > 1 \right\}.$$

Using the property of Π , we have

$$\begin{aligned} \frac{d \int \mathbb{P}_\theta d\Pi(\theta)}{d\mathbb{P}_\mu} &= \int \frac{d\mathbb{P}_\theta}{d\mathbb{P}_\mu} d\Pi(\theta) \\ &= \int \exp\left(-\frac{n}{2}\|\theta - \mu\|^2 + n\langle X - \mu, \theta - \mu \rangle\right) d\Pi(\theta) \\ &= e^{-\delta^2/2} \int \exp(n\langle X - \mu, \theta - \mu \rangle) d\Pi(\theta). \end{aligned}$$

Let $\bar{\Pi}$ be the uniform distribution on the unit sphere $\{\theta : \|\theta\| = 1\}$, and then we have

$$\int \exp(n\langle X - \mu, \theta - \mu \rangle) d\Pi(\theta) = \int \exp(\delta\sqrt{n}\langle X - \mu, \theta \rangle) d\bar{\Pi}(\theta).$$

Let f be the marginal density of the first coordinate of $\theta \sim \bar{\Pi}$. Then, $f(t) \propto (1 - t^2)^{\frac{k-3}{2}}$. The uniformity of $\bar{\Pi}$ implies that

$$\int \exp(\delta\sqrt{n}\langle X - \mu, \theta \rangle) d\bar{\Pi}(\theta) = \frac{\int_{-1}^1 \exp(\delta\sqrt{n}\|X - \mu\|t) (1 - t^2)^{\frac{k-3}{2}} dt}{\int_{-1}^1 (1 - t^2)^{\frac{k-3}{2}} dt}. \quad (56)$$

Therefore, we can write the quantity in the above display as $F(\sqrt{n}\|X - \mu\|)$. Since

$$F'(x) = \frac{\int_0^1 (e^{\delta xt} - e^{-\delta xt}) \delta t (1 - t^2)^{\frac{k-3}{2}} dt}{\int_{-1}^1 (1 - t^2)^{\frac{k-3}{2}} dt} > 0, \text{ for } x > 0,$$

the testing statistic $\frac{d \int \mathbb{P}_\theta d\Pi(\theta)}{d\mathbb{P}_\mu}$ is an increasing function of $\|X - \mu\|^2$. This implies

$$\phi = \{n\|X - \mu\|^2 \geq t\},$$

for some $t > 0$. Note that $n\|X - \mu\|^2 \sim \chi_k^2$ under \mathbb{P}_μ , and $n\|X - \mu\|^2 \sim \chi_{k,\delta^2}^2$ under any \mathbb{P}_θ with $\theta \in \bar{\Theta}_\delta$. Hence,

$$R_n(k, \delta) \geq \inf_{t>0} \left\{ \mathbb{P}(\chi_k^2 \geq t) + \mathbb{P}(\chi_{k,\delta^2}^2 < t) \right\}.$$

This completes the proof. ■

Proof [Proof of Theorem 13] Since $|\bar{\eta}_{gh}| + |\bar{\eta}_{hg}| \geq \frac{2}{|\nu_g - \nu_h|}$, it is implied by Condition M2 that $\sqrt{n}|\nu_g - \nu_h| \rightarrow \infty$ for any $g \neq h$. Moreover, for any $j \in \mathcal{C}_h$, $|\mu_j - \nu_h| = o(n^{-1/2})$. Under these assumptions, for any θ such that $\|\theta - \mu\| = \frac{\delta}{\sqrt{n}}$, there exists a $\pi \in S_k$ that depends

on θ and $\|\theta_\pi - \mu\| = \ell(\theta, \mu) = \frac{\delta}{\sqrt{n}}(1 + \epsilon_\theta)$. Moreover, $|\epsilon_\theta| = o(1)$ uniformly over all θ that satisfies $\|\theta - \mu\| = \frac{\delta}{\sqrt{n}}$. Define

$$\theta' = \mu + \frac{1}{1 + \epsilon_\theta}(\theta - \mu). \quad (57)$$

Then, $\|\theta' - \mu\| = \frac{\delta_\theta}{\sqrt{n}}$ and $\ell(\theta', \mu) = \frac{\delta}{\sqrt{n}}$, where $\delta_\theta = \frac{\delta}{1 + \epsilon_\theta}$. We use the notation R to denote the operator $R : \theta \mapsto R(\theta) = \theta'$ defined by (57). By the definition, a useful property is $\frac{R(\theta) - \mu}{\|R(\theta) - \mu\|} = \frac{\theta - \mu}{\|\theta - \mu\|}$. Consider the set

$$\bar{\Theta}_\delta = \left\{ R(\theta) : \|\theta - \mu\| = \frac{\delta}{\sqrt{n}} \right\}.$$

This definition immediately implies $\bar{\Theta}_\delta \subset \Theta_\delta$. Note that each element in $\bar{\Theta}_\delta$ can be represented as

$$R(\theta) = \mu + \frac{\delta_\theta}{\sqrt{n}} \frac{\theta - \mu}{\|\theta - \mu\|}.$$

Since there is a one-to-one relation between $\frac{\theta - \mu}{\|\theta - \mu\|}$ and a unit vector v , we can also write each element in $\bar{\Theta}_\delta$ as $\mu + \frac{\delta_\theta}{\sqrt{n}}v$. Consider a uniform probability measure $\bar{\Pi}$ on $\{v : \|v\| = 1\}$. Then, by the same argument in the proof of Theorem 11,

$$R_n(k, \delta) \geq \inf_{0 \leq \phi \leq 1} \left\{ \mathbb{P}_\mu \phi + \int \mathbb{P}_{\mu + \frac{\delta_\theta}{\sqrt{n}}v} (1 - \phi) d\bar{\Pi}(v) \right\},$$

and the likelihood ratio is

$$\mathcal{L} = \int \frac{d\mathbb{P}_{\mu + \frac{\delta_\theta}{\sqrt{n}}v}}{d\mathbb{P}_\mu} d\bar{\Pi}(v) = \int \exp(-\delta_\theta^2/2 + \delta_\theta \sqrt{n} \langle X - \mu, v \rangle) d\bar{\Pi}(v).$$

Under the assumption, there exist δ_- and δ_+ such that $\delta_- \leq \delta_\theta \leq \delta_+$ for all v and $\delta_-/\delta = 1 + o(1)$ and $\delta_+/\delta = 1 + o(1)$. We introduce the upper and lower brackets of \mathcal{L} as

$$\begin{aligned} \mathcal{L}_- &= \min \left\{ \int \exp(-\delta_+^2/2 + \delta_- \sqrt{n} \langle X - \mu, v \rangle) d\bar{\Pi}(v), \right. \\ &\quad \left. \int \exp(-\delta_+^2/2 + \delta_+ \sqrt{n} \langle X - \mu, v \rangle) d\bar{\Pi}(v) \right\}, \\ \mathcal{L}_+ &= \min \left\{ \int \exp(-\delta_-^2/2 + \delta_- \sqrt{n} \langle X - \mu, v \rangle) d\bar{\Pi}(v), \right. \\ &\quad \left. \int \exp(-\delta_-^2/2 + \delta_+ \sqrt{n} \langle X - \mu, v \rangle) d\bar{\Pi}(v) \right\}. \end{aligned}$$

The definitions imply $\mathcal{L}_- \leq \mathcal{L} \leq \mathcal{L}_+$. Define the function

$$F_\delta(x) = \frac{\int_{-1}^1 \exp(\delta xt) (1 - t^2)^{\frac{k-3}{2}} dt}{\int_{-1}^1 (1 - t^2)^{\frac{k-3}{2}} dt}. \quad (58)$$

By (56), we have

$$\begin{aligned}\mathcal{L}_- &= e^{-\delta_+^2/2} \min\{F_{\delta_-}(\sqrt{n}\|X - \mu\|), F_{\delta_+}(\sqrt{n}\|X - \mu\|)\}, \\ \mathcal{L}_+ &= e^{-\delta_-^2/2} \max\{F_{\delta_-}(\sqrt{n}\|X - \mu\|), F_{\delta_+}(\sqrt{n}\|X - \mu\|)\}.\end{aligned}$$

Define $\phi = \mathbb{I}\{\mathcal{L} > 1\}$, $\phi_- = \mathbb{I}\{\mathcal{L}_- > 1\}$ and $\phi_+ = \mathbb{I}\{\mathcal{L}_+ > 1\}$. We have the inequality $\phi_- \leq \phi \leq \phi_+$. For $\theta = \mathbb{E}X = \mu$, $\|\sqrt{n}(X - \mu)\|^2 \sim \chi_k^2$. Thus, let $Z \sim N(0, I_k)$, and then we have

$$\begin{aligned}\mathbb{P}_\mu \phi &\geq \mathbb{P}_\mu (\mathcal{L}_- > 1) \\ &= \mathbb{P}\left(e^{-\delta_+^2/2} \min\{F_{\delta_-}(\|Z\|), F_{\delta_+}(\|Z\|)\} > 1\right) \\ &\rightarrow \mathbb{P}\left(e^{-\delta^2/2} F_\delta(\|Z\|) > 1\right).\end{aligned}$$

For the alternative $\theta = \mu + \frac{\delta_v}{\sqrt{n}}v \in \bar{\Theta}_\delta$, $\|\sqrt{n}(X - \mu)\|^2 \sim \chi_{k^2, \delta_v^2}$, where $\delta_v^2 \in [\delta_-, \delta_+]$. Then,

$$\begin{aligned}\mathbb{P}_{\mu + \frac{\delta_v}{\sqrt{n}}v}(1 - \phi) &\geq \mathbb{P}_{\mu + \frac{\delta_v}{\sqrt{n}}v}(\mathcal{L}_+ \leq 1) \\ &= \mathbb{P}\left(e^{-\delta^2/2} \max\{F_{\delta_-}(\|Z + \delta_v v\|), F_{\delta_+}(\|Z + \delta_v v\|)\} \leq 1\right) \\ &\geq \mathbb{P}\left(e^{-\delta^2/2} \max\{F_{\delta_-}(\|Z + \delta_+ v\|), F_{\delta_+}(\|Z + \delta_+ v\|)\} \leq 1\right) \\ &\rightarrow \mathbb{P}\left(e^{-\delta^2/2} F_\delta(\|Z + \delta v\|) \leq 1\right)\end{aligned}$$

Note that $\mathbb{P}\left(e^{-\delta^2/2} F_\delta(\|Z + \delta v\|) \leq 1\right)$ is independent of v . Therefore, by the fact that $F_\delta(x)$ is increasing on $x > 0$, we have

$$\begin{aligned}R_n(k, n) &\geq \mathbb{P}_\mu(\mathcal{L}_- > 1) + \inf_{\|v\|=1} \mathbb{P}_{\mu + \frac{\delta_v}{\sqrt{n}}v}(\mathcal{L}_+ \leq 1) \\ &\geq (1 + o(1)) \left\{ \mathbb{P}\left(e^{-\delta^2/2} F_\delta(\|Z\|) > 1\right) + \inf_{\|v\|=1} \mathbb{P}\left(e^{-\delta^2/2} F_\delta(\|Z + \delta v\|) \leq 1\right) \right\} \\ &\geq (1 + o(1)) \inf_{t>0} \left\{ \mathbb{P}(\chi_k^2 \geq t) + \mathbb{P}(\chi_{k, \delta^2}^2 < t) \right\}.\end{aligned}$$

The proof is complete. ■

Proof [Proof of Theorem 15] Note that Condition M3 implies $\sqrt{n}|\sqrt{q_j} - \sqrt{q_l}| \rightarrow \infty$ for any $j \neq l$. Consider the set

$$\bar{\mathcal{P}}_\delta = \left\{ p : \|\sqrt{p} - \sqrt{q}\| = \frac{\delta}{\sqrt{n}} \right\}.$$

For each $p \in \bar{\mathcal{P}}_\delta$, $|\sqrt{p_j} - \sqrt{q_j}|^2 \leq \frac{\delta^2}{n}$, which implies $\sqrt{q_j}$ is the closest element to $\sqrt{p_j}$ in the set $\{\sqrt{q_1}, \dots, \sqrt{q_k}\}$. Therefore, $\ell(p, q) = \|\sqrt{p} - \sqrt{q}\| = \delta/\sqrt{n}$, which implies $\bar{\mathcal{P}}_\delta \subset \mathcal{P}_\delta$. This gives the lower bound

$$R_n(k, \delta) \geq \inf_{0 \leq \phi \leq 1} \left\{ \mathbb{P}_q \phi + \sup_{p \in \bar{\mathcal{P}}_\delta} \mathbb{P}_p(1 - \phi) \right\}.$$

Let Π be the uniform distribution on the sphere $\{v : \|v - \sqrt{q}\| = \delta/\sqrt{n}\}$. Then,

$$R_n(k, \delta) \geq \inf_{0 \leq \phi \leq 1} \left\{ \mathbb{P}_q \phi + \int \mathbb{P}_p (1 - \phi) d\Pi(\sqrt{p}) \right\}.$$

By Neyman-Pearson lemma, the optimal testing function ϕ is given by

$$\phi = \left\{ \frac{d \int \mathbb{P}_p d\Pi(\sqrt{p})}{d\mathbb{P}_q} > 1 \right\}.$$

By the definition, we have

$$\mathcal{L} = \frac{d \int \mathbb{P}_p d\Pi(\sqrt{p})}{d\mathbb{P}_q} = \int \exp \left(n \sum_{j=1}^k q_j \log \frac{p_j}{q_j} + n \sum_{j=1}^k (\hat{p}_j - q_j) \log \frac{p_j}{q_j} \right) d\Pi(\sqrt{p}),$$

where $\hat{p}_j = \frac{1}{n} \sum_{i=1}^n \mathbb{I}\{X_i = j\}$. Note that $\log \sqrt{\frac{p_j}{q_j}} = \log \left(1 + \frac{\sqrt{p_j} - \sqrt{q_j}}{\sqrt{q_j}} \right)$. By Condition M3, $\max_{1 \leq j \leq k} \left| \frac{\sqrt{p_j} - \sqrt{q_j}}{\sqrt{q_j}} \right| = o(1)$. Therefore,

$$\max_{1 \leq j \leq k} \frac{\left| \log \sqrt{\frac{p_j}{q_j}} - \frac{\sqrt{p_j} - \sqrt{q_j}}{\sqrt{q_j}} \right|}{\left| \frac{\sqrt{p_j} - \sqrt{q_j}}{\sqrt{q_j}} \right|^2} = O(1), \quad (59)$$

and

$$\max_{1 \leq j \leq k} \frac{\left| \log \sqrt{\frac{p_j}{q_j}} - \frac{\sqrt{p_j} - \sqrt{q_j}}{\sqrt{q_j}} + \frac{1}{2} \left(\frac{\sqrt{p_j} - \sqrt{q_j}}{\sqrt{q_j}} \right)^2 \right|}{\left| \frac{\sqrt{p_j} - \sqrt{q_j}}{\sqrt{q_j}} \right|^3} = O(1). \quad (60)$$

Since

$$\sum_{j=1}^k q_j \left[\frac{\sqrt{p_j} - \sqrt{q_j}}{\sqrt{q_j}} - \frac{1}{2} \left(\frac{\sqrt{p_j} - \sqrt{q_j}}{\sqrt{q_j}} \right)^2 \right] = -\|\sqrt{p} - \sqrt{q}\|^2.$$

By (60), we have

$$\sum_{j=1}^k q_j \log \frac{p_j}{q_j} = -(1 + o(1)) 2 \|\sqrt{p} - \sqrt{q}\|^2.$$

Under Condition M3, $\hat{p}_j/p_j = 1 + o_P(1)$, and this implies $\hat{p}_j/q_j = 1 + o_P(1)$. Therefore,

$$\sum_{j=1}^k (\hat{p}_j - q_j) \frac{\sqrt{p_j} - \sqrt{q_j}}{\sqrt{q_j}} = 2(1 + o_P(1)) \sum_{j=1}^k (\sqrt{\hat{p}_j} - \sqrt{q_j})(\sqrt{p_j} - \sqrt{q_j}). \quad (61)$$

By (59), we have

$$\sum_{j=1}^k (\hat{p}_j - q_j) \log \frac{p_j}{q_j} = 4(1 + o_P(1)) \sum_{j=1}^k (\sqrt{\hat{p}_j} - \sqrt{q_j})(\sqrt{p_j} - \sqrt{q_j}). \quad (62)$$

The approximations (61) and (62) imply the existence of δ_- and δ_+ that satisfies $\delta_- = (1 + o(1))\delta$, $\delta_+ = (1 + o(1))\delta$. Moreover, on an event E with probability $1 - o(1)$ under both null and alternative, the following inequalities hold:

$$-2\delta_+^2 \leq n \sum_{j=1}^k q_j \log \frac{p_j}{q_j} \leq -2\delta_-^2,$$

$$\sum_{j=1}^k (\hat{p}_j - q_j) \log \frac{p_j}{q_j} \leq \max \left\{ \frac{4\delta_-}{\sqrt{n}} \left\langle \sqrt{\hat{p}} - \sqrt{q}, \frac{\sqrt{p} - \sqrt{q}}{\|\sqrt{p} - \sqrt{q}\|} \right\rangle, \frac{4\delta_+}{\sqrt{n}} \left\langle \sqrt{\hat{p}} - \sqrt{q}, \frac{\sqrt{p} - \sqrt{q}}{\|\sqrt{p} - \sqrt{q}\|} \right\rangle \right\},$$

and

$$\sum_{j=1}^k (\hat{p}_j - q_j) \log \frac{p_j}{q_j} \geq \min \left\{ \frac{4\delta_-}{\sqrt{n}} \left\langle \sqrt{\hat{p}} - \sqrt{q}, \frac{\sqrt{p} - \sqrt{q}}{\|\sqrt{p} - \sqrt{q}\|} \right\rangle, \frac{4\delta_+}{\sqrt{n}} \left\langle \sqrt{\hat{p}} - \sqrt{q}, \frac{\sqrt{p} - \sqrt{q}}{\|\sqrt{p} - \sqrt{q}\|} \right\rangle \right\}.$$

We introduce the upper and lower brackets of the \mathcal{L} as

$$\begin{aligned} \mathcal{L}_- &= \min \left\{ \int \exp \left(-2\delta_+^2 + 4\delta_- \sqrt{n} \left\langle \sqrt{\hat{p}} - \sqrt{q}, v \right\rangle \right) d\bar{\Pi}(v), \right. \\ &\quad \left. \int \exp \left(-2\delta_+^2 + 4\delta_+ \sqrt{n} \left\langle \sqrt{\hat{p}} - \sqrt{q}, v \right\rangle \right) d\bar{\Pi}(v) \right\}, \end{aligned} \quad (63)$$

$$\begin{aligned} \mathcal{L}_+ &= \max \left\{ \int \exp \left(-2\delta_-^2 + 4\delta_- \sqrt{n} \left\langle \sqrt{\hat{p}} - \sqrt{q}, v \right\rangle \right) d\bar{\Pi}(v), \right. \\ &\quad \left. \int \exp \left(-2\delta_-^2 + 4\delta_+ \sqrt{n} \left\langle \sqrt{\hat{p}} - \sqrt{q}, v \right\rangle \right) d\bar{\Pi}(v) \right\}, \end{aligned} \quad (64)$$

where $\bar{\Pi}$ is the uniform distribution on the unit sphere $\{v : \|v\| = 1\}$. By (56), we have

$$\begin{aligned} \mathcal{L}_- &= e^{-2\delta_+^2} \min\{F_{2\delta_-}(2\sqrt{n}\|\sqrt{\hat{p}} - \sqrt{q}\|), F_{2\delta_+}(2\sqrt{n}\|\sqrt{\hat{p}} - \sqrt{q}\|)\}, \\ \mathcal{L}_+ &= e^{-2\delta_-^2} \max\{F_{2\delta_-}(2\sqrt{n}\|\sqrt{\hat{p}} - \sqrt{q}\|), F_{2\delta_+}(2\sqrt{n}\|\sqrt{\hat{p}} - \sqrt{q}\|)\}. \end{aligned}$$

where $F_\delta(x)$ is defined in (58). Note that $4n\|\sqrt{\hat{p}} - \sqrt{q}\|^2 \rightsquigarrow \chi_{k-1}^2$ under the null and $4n\|\sqrt{\hat{p}} - \sqrt{q}\|^2 - \delta_2^2 \rightsquigarrow \chi_{k-1, \delta_1^2}^2$, with $\delta_1^2 = \delta_1(p)^2$ and $\delta_2^2 = \delta_2(p)^2$ defined in (47) and (48), under the alternative. Define $\phi = \mathbb{I}\{\mathcal{L} > 1\}$, $\phi_- = \mathbb{I}\{\mathcal{L}_- > 1\}$, $\phi_+ = \mathbb{I}\{\mathcal{L}_+ > 1\}$ and $\phi^* = \mathbb{I}\{\mathcal{L}^* > 1\}$. Then, we have the inequality $\phi_- \mathbb{I}_E \leq \phi \mathbb{I}_E \leq \phi_+ \mathbb{I}_E$. For $q = p$, we have

$$\begin{aligned} \mathbb{P}_q \phi &= \mathbb{P}_q \phi \mathbb{I}_E + \mathbb{P}_q \phi \mathbb{I}_{E^c} \\ &\geq \mathbb{P}_q \phi_- \mathbb{I}_E \\ &\geq \mathbb{P}_q (\mathcal{L}_- > 1) - \mathbb{P}_q (E^c) \\ &= \mathbb{P}_q \left(e^{-2\delta_+^2} \min\{F_{2\delta_-}(2\sqrt{n}\|\sqrt{\hat{p}} - \sqrt{q}\|), F_{2\delta_+}(2\sqrt{n}\|\sqrt{\hat{p}} - \sqrt{q}\|)\} > 1 \right) - \mathbb{P}_q (E^c) \\ &\rightarrow \mathbb{P} \left(e^{-2\delta^2} F_{2\delta} \left(\sqrt{\chi_{k-1}^2} \right) > 1 \right). \end{aligned}$$

For the alternative, we have

$$\begin{aligned}
 \mathbb{P}_p(1 - \phi) &= \mathbb{P}_p(1 - \phi)\mathbb{I}_E + \mathbb{P}_p(1 - \phi)\mathbb{I}_{E^c} \\
 &\leq \mathbb{P}_p(1 - \phi_+)\mathbb{I}_E + \mathbb{P}_p(E^c) \\
 &\leq \mathbb{P}_p(1 - \phi_+) + \mathbb{P}_p(E^c) \\
 &= \mathbb{P}_p\left(e^{-2\delta^2} \max\{F_{2\delta_-}(2\sqrt{n}\|\sqrt{\hat{p}} - \sqrt{q}\|), F_{2\delta_+}(2\sqrt{n}\|\sqrt{\hat{p}} - \sqrt{q}\|)\} \leq 1\right) + \mathbb{P}_p(E^c) \\
 &\rightarrow \mathbb{P}\left(e^{-2\delta^2} F_{2\delta}\left(\sqrt{\chi_{k-1, \delta_1(p)}^2 + \delta_2(p)^2}\right) \leq 1\right) \\
 &\geq \inf_{\{\delta_1, \delta_2: \delta_1^2 + \delta_2^2 = \delta^2\}} \mathbb{P}\left(e^{-2\delta^2} F_{2\delta}\left(\sqrt{\chi_{k-1, \delta_1}^2 + \delta_2^2}\right) \leq 1\right).
 \end{aligned}$$

By the fact that $F_\delta(x)$ is increasing on $x > 0$, we have

$$\begin{aligned}
 R_n(k, \delta) &\geq (1 + o(1)) \left\{ \mathbb{P}\left(e^{-2\delta^2} F_{2\delta}\left(\sqrt{\chi_{k-1}^2}\right) > 1\right) + \right. \\
 &\quad \left. \inf_{\{\delta_1, \delta_2: \delta_1^2 + \delta_2^2 = \delta^2\}} \mathbb{P}\left(e^{-2\delta^2} F_{2\delta}\left(\sqrt{\chi_{k-1, \delta_1}^2 + \delta_2^2}\right) \leq 1\right) \right\} \\
 &\geq (1 + o(1)) \inf_{t>0} \left(\mathbb{P}\left(\chi_{k-1}^2 > t\right) + \inf_{\{\delta_1, \delta_2: \delta_1^2 + \delta_2^2 = \delta^2\}} \mathbb{P}\left(\chi_{k-1, \delta_1}^2 + \delta_2^2 \leq t\right) \right).
 \end{aligned}$$

The proof is complete. \blacksquare

Proof [Proof of Theorem 17] It is implied by Condition M4 that $\sqrt{n}|\sqrt{r_g} - \sqrt{r_h}| \rightarrow \infty$ for any $g \neq h$. Moreover, for any $j \in \mathcal{C}_h$, $|\sqrt{q_j} - \sqrt{r_h}| = o(n^{-1/2})$. Under these assumptions, for any p such that $\|\sqrt{\hat{p}} - \sqrt{q}\| = \frac{\delta}{\sqrt{n}}$, there exists a $\pi \in S_k$ that depends on p and $\|\sqrt{\hat{p}_\pi} - \sqrt{q}\| = \ell(p, q) = \frac{\delta}{\sqrt{n}}(1 + \epsilon_\theta)$. Moreover, $|\epsilon_\theta| = o(1)$ uniformly over all p that satisfies $\|\sqrt{\hat{p}} - \sqrt{q}\| = \frac{\delta}{\sqrt{n}}$. Define

$$p' = \left(\sqrt{q} + \frac{1}{1 + \epsilon_\theta} (\sqrt{\hat{p}} - \sqrt{q}) \right)^2. \tag{65}$$

Then, $\|\sqrt{\hat{p}} - \sqrt{q}\| = \frac{\delta_\theta}{\sqrt{n}}$ and $\ell(p', q) = \frac{\delta}{\sqrt{n}}$, where $\delta_\theta = \frac{\delta}{1 + \epsilon_\theta}$. We use the notation R to denote the operator $R : p \mapsto R(p)$ defined by (65). By the definition, a useful property is $\frac{\sqrt{R(p)} - \sqrt{q}}{\|\sqrt{R(p)} - \sqrt{q}\|} = \frac{\sqrt{\hat{p}} - \sqrt{q}}{\|\sqrt{\hat{p}} - \sqrt{q}\|}$. Consider the set

$$\bar{\mathcal{P}}_\delta = \left\{ R(p) : \|\sqrt{\hat{p}} - \sqrt{q}\| = \frac{\delta}{\sqrt{n}} \right\}.$$

This definition immediately implies $\bar{\mathcal{P}}_\delta \subset \mathcal{P}_\delta$. Note that each element in $\bar{\mathcal{P}}_\delta$ can be represented as

$$R(p) = \left(\sqrt{q} + \frac{\delta_\theta}{\sqrt{n}} \frac{\sqrt{\hat{p}} - \sqrt{q}}{\|\sqrt{\hat{p}} - \sqrt{q}\|} \right)^2.$$

Since there is a one-to-one relation between $\frac{\sqrt{p}-\sqrt{q}}{\|\sqrt{p}-\sqrt{q}\|}$ and a unit vector v , we can also write each element in $\bar{\mathcal{P}}_\delta$ as $(\sqrt{q} + \frac{\delta_v}{\sqrt{n}}v)^2$. Consider a uniform probability measure $\bar{\Pi}$ on $\{v : \|v\| = 1\}$. Then, by the same argument in the proof of Theorem 11,

$$R_n(k, \delta) \geq \inf_{0 \leq \phi \leq 1} \left\{ \mathbb{P}_q \phi + \int \mathbb{P} \left(\sqrt{q} + \frac{\delta_v}{\sqrt{n}}v \right)^2 (1 - \phi) d\bar{\Pi}(v) \right\},$$

and the likelihood ratio is

$$\mathcal{L} = \int \frac{d\mathbb{P} \left(\sqrt{q} + \frac{\delta_v}{\sqrt{n}}v \right)^2}{d\mathbb{P}_q} d\bar{\Pi}(v).$$

Using the same arguments in the proofs of Theorem 13 and Theorem 15, there exist δ_- and δ_+ , with which we can define \mathcal{L}_- and \mathcal{L}_+ as in (63) and (64) with the desired properties. Then, the same argument in the proof of Theorem 15 leads to the desired result. ■

Proof [Proof of Theorem 12] By studying the proof of Theorem 22, the only probabilistic argument in approximation is that $\max_{1 \leq j \leq k} Z_j^2 \leq C_n$ in probability. Since this event is independent of θ , the in-probability argument can be made uniformly over $\theta \in \Theta_\delta$ and $\theta \in \Theta_0$. ■

Proof [Proof of Theorem 14] By Theorem 23, $T_g \geq T_f$ in probability. This implies that $\mathbb{P}_\theta \phi = \mathbb{P}_\theta(T_g > t^*)$ and $\mathbb{P}_\theta(1 - \phi) = \mathbb{P}_\theta(T_g \leq t^*)$ under both null and alternative distributions. Then, by the same argument in the proof of Theorem 12, we obtain the desired conclusion. ■

Proof [Proofs of Theorem 16 and Theorem 18] Similar to the argument used in the proof of Theorem 12, the results directly follow the conclusions of Theorem 24 and Theorem 25. ■

Acknowledgments

The research of CG is supported in part by NSF grant DMS-1712957.

References

- Animashree Anandkumar, Daniel Hsu, and Sham M Kakade. A method of moments for mixture models and hidden markov models. In *Conference on Learning Theory*, pages 33–1, 2012.
- Animashree Anandkumar, Rong Ge, Daniel Hsu, and Sham Kakade. A tensor spectral approach to learning mixed membership community models. In *Conference on Learning Theory*, pages 867–881, 2013.

- Francis J Anscombe. The transformation of poisson, binomial and negative-binomial data. *Biometrika*, 35(3/4):246–254, 1948.
- Forough Arabshahi and Animashree Anandkumar. Spectral methods for correlated topic models. *arXiv preprint arXiv:1605.09080*, 2016.
- Sanjeev Arora, Rong Ge, and Ankur Moitra. New algorithms for learning incoherent and overcomplete dictionaries. In *Conference on Learning Theory*, pages 779–806, 2014.
- Debapratim Banerjee and Zongming Ma. Optimal hypothesis testing for stochastic block models with growing degrees. *arXiv preprint arXiv:1705.05305*, 2017.
- Gilles Blanchard, Alexandra Carpentier, and Maurilio Gutzeit. Minimax euclidean separation rates for testing convex hypotheses in \mathbb{R}^d . *arXiv preprint arXiv:1702.03760*, 2017.
- Harry Crane. The ubiquitous ewens sampling formula. *Statistical Science*, 31(1):1–19, 2016.
- Pierpaolo De Blasi, Stefano Favaro, Antonio Lijoi, Ramsés H Mena, Igor Prünster, and Matteo Ruggiero. Are gibbs-type priors the most natural generalization of the dirichlet process? *IEEE transactions on pattern analysis and machine intelligence*, 37(2):212–229, 2015.
- Warren J Ewens. The sampling theory of selectively neutral alleles. *Theoretical population biology*, 3(1):87–112, 1972.
- Chao Gao and John Lafferty. Testing network structure using relations between small subgraph probabilities. *arXiv preprint arXiv:1704.06742*, 2017.
- Alexander Gnedin and Jim Pitman. Exchangeable gibbs partitions and stirling triangles. *Journal of Mathematical sciences*, 138(3):5674–5685, 2006.
- Roger A Horn and Charles R Johnson. *Matrix analysis*. Cambridge university press, 2012.
- Daniel Hsu and Sham M Kakade. Learning mixtures of spherical gaussians: moment methods and spectral decompositions. In *Proceedings of the 4th conference on Innovations in Theoretical Computer Science*, pages 11–20. ACM, 2013.
- John FC Kingman. Random discrete distributions. *Journal of the Royal Statistical Society. Series B (Methodological)*, pages 1–22, 1975.
- S Kocherlakota and K Kocherlakota. Goodness of fit tests for discrete distributions. *Communications in statistics-theory and methods*, 15(3):815–829, 1986.
- RJ Kulperger and AC Singh. On random grouping in goodness of fit tests of discrete distributions. *Journal of Statistical Planning and Inference*, 7(2):109–115, 1982.
- Peter Müller, Abel Rodriguez, et al. Random partition models. In *Nonparametric Bayesian Inference*, pages 87–92. IMS and ASA, 2013.

Karl Pearson. On the criterion that a given system of deviations from the probable in the case of a correlated system of variables is such that it can be reasonably supposed to have arisen from random sampling. *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science*, 50(302):157–175, 1900.

Amelia Perry, Jonathan Weed, Afonso Bandeira, Philippe Rigollet, and Amit Singer. The sample complexity of multi-reference alignment. *arXiv preprint arXiv:1707.00943*, 2017.

Jim Pitman. Exchangeable and partially exchangeable random partitions. *Probability theory and related fields*, 102(2):145–158, 1995.

Jim Pitman. Random discrete distributions invariant under size-biased permutation. *Advances in Applied Probability*, 28(2):525–539, 1996.